antibodies -online.com





anti-HCN1 antibody (AA 778-910) (Atto 594)





Go to Product page

\sim				
()\/	Δr	1//		۱۸
Οv	CI	۷I	て	V١

Quantity:	100 μg
Target:	HCN1
Binding Specificity:	AA 778-910
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HCN1 antibody is conjugated to Atto 594
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Antibody Array (AA)

Product Details

Immunogen:	Fusion protein amino acids 778-910 (C terminus) of rat HCN1
Clone:	S70
Isotype:	IgG1
Specificity:	Detects ~100 kDa. No cross-reactivity against HCN2.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details


HCN1

Target Details

Storage Comment:

l arget Details	
Alternative Name:	HCN1 (HCN1 Products)
Background:	Hyperpolarization-activated cation channels of the HCN gene family, such as HCN1, play a crucial role in the regulatons of cell excitability. Importantly, they contribute to spontaneous rhythmic activity in both the heart and brain (1).
Gene ID:	84390
NCBI Accession:	NP_445827
UniProt:	Q9JKB0
Pathways:	Asymmetric Protein Localization
Application Details	
Application Notes:	 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2483957 was sufficient for detection of HCN1 in 10 μg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
_	

Conjugated antibodies should be stored at 4°C








Image 2. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

Image 1. Western Blot analysis of Rat brain membrane

lysate showing detection of HCN1 protein using Mouse Anti-

HCN1 Monoclonal Antibody, Clone S70-28 . Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody:

Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 2

hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG:



Immunohistochemistry

Western Blotting

HRP for 1 hour at RT.

Image 3. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Tissue: Frozen brain section. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 μl for 5 minutes at RT. Localization: Neurons.

Please check the product details page for more images. Overall 5 images are available for ABIN2483957.