

Datasheet for ABIN2483963
anti-HCN1 antibody (AA 778-910) (APC)

5 Images

[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HCN1
Binding Specificity:	AA 778-910
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HCN1 antibody is conjugated to APC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Antibody Array (AA)

Product Details

Immunogen:	Fusion protein amino acids 778-910 (C terminus) of rat HCN1
Clone:	S70
Isotype:	IgG1
Specificity:	Detects ~100 kDa. No cross-reactivity against HCN2.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

Target:	HCN1
---------	------

Target Details


Alternative Name:	HCN1 (HCN1 Products)
Background:	Hyperpolarization-activated cation channels of the HCN gene family, such as HCN1, play a crucial role in the regulations of cell excitability. Importantly, they contribute to spontaneous rhythmic activity in both the heart and brain (1).
Gene ID:	84390
NCBI Accession:	NP_445827
UniProt:	Q9JKB0
Pathways:	Asymmetric Protein Localization

Application Details

Application Notes:	<ul style="list-style-type: none">• WB (1:1000)• IHC (1:1000)• ICC/IF (1:100)• optimal dilutions for assays should be determined by the user.
Comment:	1 µg/ml of ABIN2483963 was sufficient for detection of HCN1 in 10 µg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only


Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C




Immunohistochemistry

Image 1. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Tissue: Frozen brain section. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT. Localization: Neurons.



Immunohistochemistry

Image 2. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Tissue: Frozen brain section. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-HCN1 Monoclonal Antibody, Clone S70-28 . Tissue: Cerebellum. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-HCN1 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 250-500 µl for 5 minutes at RT. Localization: Cytoplasmic staining of

Purkinje cells.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN2483963.