antibodies .- online.com







anti-MDC1 antibody (N-Term) (HRP)







()	1 /	-	r٧	/ 1	0	A .
	1//	\vdash	1 \/	/ I	_	۱/۱
\sim	٧.	\sim	1 V		$\overline{}$	V١

Quantity:	100 μg	
Target:	MDC1	
Binding Specificity:	N-Term	
Reactivity:	Mouse	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This MDC1 antibody is conjugated to HRP	
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)	

Product Details


Immunogen:	GST-tagged recombinant protein corresponding to mouse MDC1 at and around the N-terminus	
Clone:	P2B11	
Isotype:	lgG1	
Specificity:	Detects ~184 kDa. This antibody recognizes MDC1 at and around the N-terminus.	
Cross-Reactivity:	Chimpanzee, Cow, Human, Mouse	
Purification:	Protein G Purified	

Target Details


Target:	MDC1
Alternative Name:	MDC1 (MDC1 Products)

Target Details

rarget Details		
Background:	MDC1, mediator of DNA damage checkpoint protein 1, plays a role in checkpoint mediated cell cycle arrest in response to DNA damage, within S phase and G2/M. It is also thought to act as a scaffold protein during recruitment of DNA repair and signal transduction proteins to discrete foci of DNA damage that are marked by phosphorylation of histone H2A.X on S139.	
Gene ID:	240087	
NCBI Accession:	NP_001010833	
UniProt:	Q5PSV9	
Application Details		
Application Notes:	 WB (1:2000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user. 	
Comment:	$0.5\mu g/ml$ of ABIN2484153 was sufficient for detection of MDC1 in 10 μg of HeLa cell lysate by ECL immunoblot analysis.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Conjugated antibodies should be stored at 4°C	



kDa MW 2 250 150 100 75



Immunofluorescence (fixed cells)

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-MDC1 Monoclonal Antibody, Clone P2B11. Tissue: Fibroblast cell line (NIH 3T3). Species: Mouse. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-MDC1 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: DAPI (blue) nuclear stain at 1:5000 for 5 min RT. Localization: Nucleus. Magnification: 60X.

Western Blotting

Image 2. Western Blot analysis of Human 293Trap cell lysates showing detection of 184 kDa MDC1 protein using Mouse Anti-MDC1 Monoclonal Antibody, Clone P2B11. Lane 1: MW ladder. Lane 2: 293Trap cell lysates. Load: 30 µg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-MDC1 Monoclonal Antibody at 1:1000 for 2 hours RT. Secondary Antibody: Goat Anti-Mouse HRP: lgG at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 184 kDa.

Western Blotting

Image 3. Western Blot analysis of Mouse Cortex and Cerebellum showing detection of 184 kDa MDC1 protein using Mouse Anti-MDC1 Monoclonal Antibody, Clone P2B11 . Lane 1: MW ladder. Lane 2: Mouse Cortex. Lane 3: Mouse Cerebellum. Load: 10 μg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-MDC1 Monoclonal Antibody at 1:1000 for 2 hours RT. Secondary Antibody: Goat Anti-Mouse at 1:2000 for 60 min at RT. Color Development: ECL solution for 5 min in RT. Predicted/Observed Size: 184 kDa.