antibodies -online.com







Images



Overview

Quantity:	200 μg
Target:	Hsc70 (HSPA8)
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Hsc70 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC), Proximity Ligation Assay (PLA), Binding Studies (Bind), Antibody Array (AA)

Product Details

Target:

1 Toddet Details		
Immunogen:	Full length human HSC70	
Clone:	1F2-H5	
Isotype:	IgG2a kappa	
Specificity:	Detects ~73 kDa. Does not cross react with HSP70.	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Protein G Purified	
Target Details		

Hsc70 (HSPA8)


Target Details

Alternative Name:	HSC70 (HSP73) (HSPA8 Products)
Background:	HSP70 genes encode abundant heat-inducible 70- kDa HSPs (HSP70s). In most eukaryotes
	HSP70 genes exist as part of a multigene family. They are found in most cellular compartments
	of eukaryotes including nuclei, mitochondria, chloroplasts, the endoplasmic reticulum and the
	cytosol, as well as in bacteria. The genes show a high degree of conservation, having at least
	50 % identity (2). The N-terminal two thirds of HSP70s are more conserved than the C-terminal
	third. HSP70 binds ATP with high affinity and possesses a weak ATPase activity which can be
	stimulated by binding to unfolded proteins and synthetic peptides (3). When HSC70
	(constitutively expressed) present in mammalian cells was truncated, ATP binding activity was
	found to reside in an N-terminal fragment of 44 kDa which lacked peptide binding capacity.
	Polypeptide binding ability therefore resided within the C-terminal half (4). The structure of this
	ATP binding domain displays multiple features of nucleotide binding proteins (5). When cells
	are subjected to metabolic stress (e.g., heat shock) a member of the HSP 70 family, HSP 70
	(HSP72), is expressed, HSP 70 is highly related to HSC70 (>90 % sequence identity).
	Constitutively expressed HSC70 rapidly forms a stable complex with the highly inducible HSP70
	in cells following heat shock. The interaction of HSC70 with HSP 70 is regulated by ATP. These
	two heat shock proteins move together in the cell experiencing stress. Furthermore, research
	on HSC70 has implicates it with a role in facilitating the recovery of centrosomal structure and
	function after heat shock (6).
Gene ID:	3312
NCBI Accession:	NP_006588
UniProt:	P11142
Application Details	
Application Notes:	• WB (1:1000)
	• ICC/IF (1:100)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μg/ml of ABIN2484322 was sufficient for detection of HSC70 in 10 μg of HeLa lysate by
	colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid

Handling


Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

Images




Immunofluorescence (fixed cells)

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Hsc70 (Hsp73) Monoclonal Antibody, Clone 1F2-H5. Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-Hsc70 (Hsp73) Monoclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Melanosome. Localizes to nucleus upon heat shock. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsc70 (Hsp73) Antibody. (C) Composite.



Western Blotting

Image 2. Western Blot analysis of Human Cell lysates showing detection of Hsc70 protein using Mouse Anti-Hsc70 Monoclonal Antibody, Clone 1F2-H5 . Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-Hsc70 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT. 1: mix of 10 different human cell lines, 2: Hsp72 recombinant protein, and 3: Hsc70(Hsp73) recombinant protein.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Hsc70 (Hsp73) Monoclonal Antibody, Clone 1F2-H5. Tissue: Heat Shocked HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Mouse Anti-Hsc70 (Hsp73) Monoclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: R-PE Goat Anti-Mouse (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Melanosome. Localizes to nucleus upon heat shock. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsc70 (Hsp73) Antibody. (C) Composite.

Please check the product details page for more images. Overall 4 images are available for ABIN2484322.