# antibodies - online.com







## anti-Bassoon antibody (AA 786-1041) (Biotin)

**Images** 



| $\sim$ |          |
|--------|----------|
| ( )\/△ | rview    |
| $\cup$ | 1 410 44 |

| Quantity:            | 100 μg  |
|----------------------|---|
| Target:              | Bassoon (BSN)                                     |
| Binding Specificity: | AA 786-1041                                       |
| Reactivity:          | Human   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This Bassoon antibody is conjugated to Biotin     |
| Application:         | Immunohistochemistry (IHC), Western Blotting (WB) |

#### **Product Details**

| Immunogen:        | NM_003458.3 (AA 786-1041) N-terminal his-tagged fusion protein |
|-------------------|--|
| Specificity:      | Detects ~420 kDa. Multiple isoforms can be detected.           |
| Cross-Reactivity: | Human, Mouse, Rat  |
| Purification:     | Protein A Purified   |

### Target Details

| Target:           | Bassoon (BSN)  |
|-------------------|--|
| Alternative Name: | Bassoon (BSN Products)   |
| Background:       | Bassoon is a 420 kDa protein that is a localized at the presynaptic nerve terminals and is           |
|                   | believed to play a role in the structural and functional organization of the synaptic vesicle cycle. |

| Bassoon is predicted to contain two double-zinc fingers, three coiled-coil regions, an  | d two          |
|---|----------------|
| polyglutamine domains. The polyglutamine domains in the C-terminus are of interes       | t, since it is |
| known that for some human proteins, such as Huntington, abnormal amplification o        | f this         |
| region can cause late-onset neurodegeneration. Bassoon is concentrated at sites op      | posite to      |
| postsynaptic densities in synaptic terminals and in cultured neurons, it is found to co | -localize      |
| with GABA (A) and glutamate (GluR1) receptors (1).                                      |                |

| Gene ID:        | 8927      |
|-----------------|-----------|
| NCBI Accession: | NP_003449 |
| UniProt:        | Q9UPA5    |

## **Application Details**

| Application Notes: | <ul><li>WB (1:1000)</li><li>IHC (1:400)</li></ul>   |
|--------------------|---|
|                    | optimal dilutions for assays should be determined by the user.  |
| Comment:           | 1 $\mu$ g/ml of ABIN2484542 was sufficient for detection of Bassoon in 10 $\mu$ g of rat brain tissue lysate by colorimetric immunoblot analysis using goat ant rabbit IgG:HRP as the secondary |
|                    | antibody.   |

For Research Use only

## Handling


Restrictions:

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 mg/mL  |
| Buffer:            | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated                              |
| Preservative:      | Sodium azide   |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage:           | 4 °C   |
| Storage Comment:   | Conjugated antibodies should be stored at 4°C  |



#### **Immunohistochemistry**

Image 1. Immunohistochemistry analysis using Rabbit Anti-Bassoon Polyclonal Antibody . Tissue: Muscle. Species: Mouse. Primary Antibody: Rabbit Anti-Bassoon Polyclonal Antibody at 1:400. Secondary Antibody: Alexa Fluor 488 Goat Anti-Rabbit. Counterstain: BTX (red). Localization: Selective staining of the NMJ.



#### **Western Blotting**

**Image 2.** Western blot analysis of Mouse, Rat brain cell lysates showing detection of Bassoon protein using Rabbit Anti-Bassoon Polyclonal Antibody . Primary Antibody: Rabbit Anti-Bassoon Polyclonal Antibody at 1:1000.