

Datasheet for ABIN2484714
anti-HDEL antibody (C-Term) (FITC)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	HDEL
Binding Specificity:	C-Term
Reactivity:	Saccharomyces cerevisiae
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This HDEL antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (IF), Immunocytochemistry (ICC)

Product Details

Immunogen:	Raised against a synthetic HDEL peptide corresponding to the C-terminus of yeast Bip
Clone:	2E7
Isotype:	IgG2b
Specificity:	Detects ~78 kDa.
Cross-Reactivity:	Drosophila melanogaster, Plant, Saccharomyces cerevisiae
Purification:	Protein G Purified

Target Details

Target:	HDEL
Alternative Name:	HDEL (HDEL Products)

Target Details

Background:	HSP 70 family comprises four highly conserved proteins, HSP 70, HSC 70, GRP 75 and GRP 78, which serve a variety of roles. They act as molecular chaperones, facilitating the assembly of multi-protein complexes, participate in the translocation of polypeptides across cell membranes and to the nucleus, and aid in the proper folding of nascent polypeptide chains (1, 2). GRP 78 is localized in the endoplasmic reticulum (ER), where it receives imported secretory proteins and is involved in the folding and translocation of nascent peptide chains (2). Sorting of these proteins is dependent on a C-terminal tetrapeptide signal, usually KDEL in animal cells, and HDEL in <i>S.cerevisiae</i> (3). The 2E7 clone recognizes the C-terminal peptide HDEL, a common version of the endoplasmic reticulum retention signal found in yeast, plant, nematode and other ER proteins. 2E7 specifically stains HDEL proteins in barnyard grass, beet, cotton, mung bean, sorghum and wheat (4).
-------------	--

Application Details

Application Notes:	<ul style="list-style-type: none">• WB (1:1000)• ICC/IF (1:100)• optimal dilutions for assays should be determined by the user.
Comment:	1 µg/ml of ABIN2484714 was sufficient for detection of HDEL-containing proteins in 10 µg of <i>S. cerevisiae</i> lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C