

Datasheet for ABIN2484822  
**anti-DLG4 antibody (Atto 390)**



[Go to Product page](#)

4 Images

### Overview

Quantity:	100 µg
Target:	DLG4
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This DLG4 antibody is conjugated to Atto 390
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC), Antibody Array (AA)

### Product Details

Immunogen:	Recombinant rat PSD-95
Clone:	7E3
Isotype:	IgG1
Specificity:	Detects ~100 kDa. An additional protein of >100 kDa is also detected. Additional cross-reactive bands are detected at ~75 kDa and 50 kDa in rat and mouse samples.
Cross-Reactivity:	Cow, Human, Mouse, Rat
Purification:	Protein G Purified

### Target Details

Target:	DLG4
---------	------

## Target Details

---

Alternative Name: PSD95 ([DLG4 Products](#))

---

Background: Postsynaptic Density protein 95 (PSD95), also known as Synapse associated protein 90 kDa, is a member of the membrane-associated guanylate kinase (MAGUK) family of proteins. PSD95 is a scaffolding protein and is involved in the assembly and function of the postsynaptic density complex (1). These family members consist of an N-terminal variable segment followed by three amino-terminal PDZ domains, an upstream SH3 domain and an inactive carboxyl-terminal guanylate kinase (GK) domain. The first and second PDZ domain localize NMDA receptors and K<sup>+</sup> channels to synapses, and the third binds to neuroligins which are neuronal cell adhesion molecules that interact with b-neurexins and form intercellular junctions. PSD-95 also binds to neuronal nitric oxide synthase, possibly through interactions between PDZ domains present on both proteins (2). Thus different PDZ domains of PSD-95 might be specialized for distinct functions (3, 4). PSD95 participates in synaptic targeting of AMPA receptors through an indirect manner involving Stargazin and related transmembrane AMPA receptor regulatory proteins (TARPs) (5). The protein is implicated in experience dependent plasticity and plays an indispensable role in learning (6). Mutations in PSD95 are associated with autism (7).

---

Gene ID: 29495

---

NCBI Accession: [NP\\_062567](#)

---

UniProt: [P31016](#)

---

Pathways: [Regulation of Muscle Cell Differentiation](#), [Synaptic Membrane](#), [Skeletal Muscle Fiber Development](#), [Asymmetric Protein Localization](#), [Regulation of long-term Neuronal Synaptic Plasticity](#)

---

## Application Details

---

Application Notes:

- WB (1:1000)
- IHC (1:1000)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

---

Comment: 1 µg/ml was sufficient for detection of PSD-95 on 20 µg rat brain tissue extract by ECL immunoblot analysis using Goat Anti-Mouse IgG: HRP as the secondary.

---

Restrictions: For Research Use only

---

## Handling

---


Format: Liquid

---

## Handling

Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

## Images




### Immunohistochemistry

**Image 1.** Immunohistochemistry analysis using Mouse Anti-PSD95 Monoclonal Antibody, Clone 7E3 . Tissue: Neocortex. Species: Rat. Primary Antibody: Mouse Anti-PSD95 Monoclonal Antibody at 1:1000.



### Immunofluorescence (fixed cells)

**Image 2.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-PSD95 Monoclonal Antibody, Clone 7E3 . Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-PSD95 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Filamentous-like staining.



### Immunohistochemistry

**Image 3.** Immunohistochemistry analysis using Mouse Anti-PSD95 Monoclonal Antibody, Clone 7E3 . Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-PSD95 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Basal cell staining in the epidermis, some hair follicle staining, dermal staining.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN2484822.