antibodies

Datasheet for ABIN2485004 anti-GABRA1 antibody (AA 15-34) (Biotin)





Overview

Quantity:	100 µg	
Target:	GABRA1	
Binding Specificity:	AA 15-34	
Reactivity:	Rat	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This GABRA1 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)	

Product Details


Immunogen:	Synthetic peptide amino acids 15-34 (N-terminus) of rat GABA-A-R-Delta
Clone:	N151-3 (Formerly S151-3)
Isotype:	lgG2a
Specificity:	Detects ~55 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified
Target Details	
Target:	GABRA1

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN2485004 | 09/11/2023 | Copyright antibodies-online. All rights reserved.


Target Details	
Alternative Name:	GABA A Receptor (GABRA1 Products)
Background:	The GABA-A receptor is a member of the superfamily of fast acting ligand-gated ion channels. The individual subunits of these receptors have similar sequences and structural features (1). GABA-A receptors are the major fast inhibitory neurotransmitter gated ion channels in the brain (2).
Gene ID:	29689
NCBI Accession:	NP_058985
UniProt:	P18506
Application Details	
Application Notes:	 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user.
Comment:	2 μg/ml of ABIN2485004 was sufficient for detection of Delta1 GABA-A receptor in 10 μg of rat brain lysate by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

Storage Comment: Conjugated antibodies should be stored at 4°C

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN2485004 | 09/11/2023 | Copyright antibodies-online. All rights reserved.



79.68→	1
48.33→	-
37.81→	and a
23.27→	
18.19→	



Immunocytochemistry

Immunocytochemistry/Immunofluorescence Image 1. analysis using Mouse Anti-GABA-A Receptor Delta Monoclonal Antibody, Clone N151/3 (ABIN2485004). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Monoclonal Delta GABA-A Receptor Antibody (ABIN2485004) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) GABA-A Receptor Delta Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Rat Cell line lysates showing detection of GABA A Receptor protein using Mouse Anti-GABA A Receptor Monoclonal Antibody, Clone S151-3 . Load: 15 µg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-GABA A Receptor Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-GABA A Receptor Monoclonal Antibody, Clone S151-3 . Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at - 20°C. Primary Antibody: Mouse Anti-GABA A Receptor Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Diffuse cytoplasm and dull nuclei.

Please check the product details page for more images. Overall 4 images are available for ABIN2485004.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN2485004 | 09/11/2023 | Copyright antibodies-online. All rights reserved.