# antibodies -online.com





## anti-ADRA2C antibody (AA 442-462) (Atto 594)

3 Images




Go to Product page


$\sim$					
	1//	⊃r	V/I	Φ\	Λ

Quantity:	100 μg	
Target:	ADRA2C	
Binding Specificity:	AA 442-462	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This ADRA2C antibody is conjugated to Atto 594	
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)	
Product Details		
Immunogen:	Synthetic peptide amino acids 442- 462 (QDFRRSFKHILFRRRRRGFRQ, cytoplasmic C-	
	terminus) of human Alpha-2C adrenergic receptor	
Clone:	S330A-80	
Isotype:	lgG1	
Specificity:	Detects 50 kDa or larger (possibly due to dimerization). Does not cross-react with other	
	adrenergic receptors.	
Cross-Reactivity:	Human, Mouse	
Purification:	Protein G Purified	

### Target Details

Target:	ADRA2C	
Alternative Name:	Alpha2C Adrenergic (ADRA2C Products)	
Background:	The alpha 2C Adrenergic Receptor controls the release of neurotransmitter from central adrenergic neurons and from sympathetic nerves in the heart. This receptor also plays a role in cognitive and behavioral function. Two variants are produced by alternative splicing. Alpha 2C Adrenergic Receptor has been reported extensively in the brain. Blood, corpus cavermosum, kidney, and heart tissues have also been reported to express the protein.	
Gene ID:	152	
NCBI Accession:	NP_000674	
UniProt:	P18825	
Pathways:	EGFR Signaling Pathway, cAMP Metabolic Process	
Application Details		
Application Notes:	<ul> <li>WB (1:1000)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>	
Comment:	$1 \mu$ g/ml of ABIN2485201 was sufficient for detection of Alpha2C Adrenergic Receptor in 20 μg of COS cells transiently transfected with HA-tagged Alpha- lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Conjugated antibodies should be stored at 4°C	






#### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 (ABIN2485201). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody (ABIN2485201) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Alpha-2C Adrenergic Receptor Antibody (D) Composite.

#### **Western Blotting**

Image 2. Western Blot analysis of Monkey COS cells transfected with HA-tagged Alpha-2C showing detection of ~50 kDa Alpha-2C Adrenergic Receptor protein using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 . Lane 1: Molecular Weight Ladder. Lane 2: Monkey COS cells transfected with HA-tagged Alpha-2C. Load: 15 μg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody at 1:200 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~50 kDa.



#### Immunofluorescence (fixed cells)

Immunocytochemistry/Immunofluorescence **Image** 3. analysis using Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody, Clone S330A-80 Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Alpha-2C Adrenergic Receptor Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cell Membrane, Nucleus. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Alpha-2C Adrenergic Receptor Antibody (D) Composite.