# antibodies -online.com







# anti-TRPV3 antibody (AA 458-474) (PerCP)



**Images** 



# Overview

Quantity:	100 μg
Target:	TRPV3
Binding Specificity:	AA 458-474
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TRPV3 antibody is conjugated to PerCP
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)


# **Product Details**


Immunogen:	Synthetic peptide amino acids 458-474 (cytoplasmic C-terminus) of rat TrpV3
Clone:	N15-39 (Formerly S15-39)
Isotype:	lgG2a
Specificity:	Detects ~70 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified


# **Target Details**

# **Target Details**

etion in a ermo sensitiv ate in the skin, d at nother family ought to
cell lysate at anti-mouse
n conjugated
ICE which
at







### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39 (ABIN2485246). Tissue: HaCaT cells. Species: Human. Fixation: Cold 100 % methanol for 10 minutes at -20 °C. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody (ABIN2485246) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Dotty staining in all cells. Some intermediate filament-like staining in some cells.

### **Western Blotting**

Image 2. Western Blot analysis of Rat brain membrane lysate showing detection of TrpV3 protein using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39 . Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

## **Immunohistochemistry**

Image 3. Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Filaggrin-like staining in upper layers. Dull lower layer cell staining. Some stain seen in hypodermis.

Please check the product details page for more images. Overall 5 images are available for ABIN2485246.