antibodies - online.com







anti-TRPV3 antibody (AA 458-474) (PE)



Images



\sim			
	IV/E	۱/۱۲	$I \cap V$

Quantity:	100 μg	
Target:	TRPV3	
Binding Specificity:	AA 458-474	
Reactivity:	Rat	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This TRPV3 antibody is conjugated to PE	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)	

Product Details


Immunogen:	Synthetic peptide amino acids 458-474 (cytoplasmic C-terminus) of rat TrpV3	
Clone:	N15-39 (Formerly S15-39)	
Isotype:	lgG2a	
Specificity:	Detects ~70 kDa.	
Cross-Reactivity:	Human, Mouse, Rat	
Purification:	Protein G Purified	

Target Details

|--|


Target Details

TRPV3 (TRPV3 Products)	
The TRPV3 protein belongs to a family of non-selective cation channels that function in a variety of processes, including temperature sensation and vasoregulation. The thermo sensitive members of this family are expressed in subsets of sensory neurons that terminate in the sking and are activated at distinct physiological temperatures. This channel is activated at temperatures between 22 and 40 degrees C. The gene lies in close proximity to another family member (TRPV1) gene on chromosome 17, and the two encoded proteins are thought to associate with each other to form heteromeric channels (1, 2).	
497948	
NP_001020928	
Q4QYD9	
 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user. 	
1 μg/ml of ABIN2485247 was sufficient for detection of TrpV3 in 10 μg of COS-1 cell lysate transiently transfected with TrpV3 by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody	
For Research Use only	
Liquid	
1 mg/mL	
PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Sodium azide	
This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
4 °C	
Conjugated antibodies should be stored at 4°C	




Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39 (ABIN2485247). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody (ABIN2485247) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) TrpV3 Antibody (D) Composite.



Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39. Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Dotty staining in all cells. Some intermediate filament-like staining in some cells.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-39. Tissue: backskin. Species: Mouse. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Filaggrin-like staining in upper layers. Dull lower layer cell staining. Some stain seen in hypodermis.

Please check the product details page for more images. Overall 5 images are available for ABIN2485247.