



[Go to Product page](#)

Datasheet for ABIN2485328

## anti-SLC18A3 antibody (AA 521-532) (FITC)

### 3 Images

#### Overview

Quantity:	100 µg
Target:	SLC18A3
Binding Specificity:	AA 521-532
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This SLC18A3 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC)

#### Product Details

Immunogen:	Synthetic peptide amino acids 521-532 of human VACHT
Clone:	S6-38
Isotype:	IgG1
Specificity:	Detects ~56 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

#### Target Details

Target:	SLC18A3
---------	---------

## Target Details

---

Alternative Name: VACHT ([SLC18A3 Products](#))

---

Background: VACHT is a member of the vesicular amine transporter (VMAT) family. The encoded transmembrane protein transports acetylcholine into secretory vesicle for release into the extracellular space. Acetylcholine (Ach) transport utilizes a proton gradient established by a vacuolar ATPase. This gene is located within the first intron of the choline acetyltransferase gene.

---

Gene ID: 6572

---

NCBI Accession: [NP\\_003046](#)

---

UniProt: [Q16572](#)

---

## Application Details

---

Application Notes:

- WB (1:1000)
- IHC (1:200)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

---

Comment: A dilution of 1:50-1:200 of SMC-341 was sufficient for detection of VACHT Transporter in rat brain using immunohistochemistry analysis and goat anti-mouse IgG:HRP as the secondary antibody.

---

Restrictions: For Research Use only

---

## Handling

---

Format: Liquid

---

Concentration: 1 mg/mL

---

Buffer: PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

---

Preservative: Sodium azide

---

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


---

Storage: 4 °C

---


Storage Comment: Conjugated antibodies should be stored at 4°C

---




### Immunocytochemistry

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VACHT Monoclonal Antibody, Clone S6-38 (ABIN2485328). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-VACHT Monoclonal Antibody (ABIN2485328) at 1:200 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) VACHT Antibody (D) Composite.



### Immunofluorescence (fixed cells)

**Image 2.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-VACHT Monoclonal Antibody, Clone S6-38 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-VACHT Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:200 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min at RT, 5 min at RT. Localization: Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) VACHT Antibody (D) Composite.



### Western Blotting

**Image 3.** Western Blot analysis of Rat brain membrane lysate showing detection of VACHT protein using Mouse Anti-VACHT Monoclonal Antibody, Clone S6-38 . Primary Antibody: Mouse Anti-VACHT Monoclonal Antibody at 1:1000.