
antibodies - online.com







anti-TRPV3 antibody (AA 774-791) (FITC)





Overview

Quantity:	100 μg
Target:	TRPV3
Binding Specificity:	AA 774-791
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This TRPV3 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)

Product Details


Immunogen:	Synthetic peptide amino acids 774-791 (C-terminus) of rat TrpV3
Clone:	N15-4 (Formerly S15-4)
Isotype:	lgG2a
Specificity:	Detects ~70 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details


|--|

Target Details

rarget Details	
Alternative Name:	TRPV3 (TRPV3 Products)
Background:	The TRPV3 protein belongs to a family of nonselective cation channels that function in a variet
	of processes, including temperature sensation and vasoregulation. The thermosensitive
	members of this family are expressed in subsets of sensory neurons that terminate in the skin,
	and are activated at distinct physiological temperatures. This channel is activated at
	temperatures between 22 and 40 degrees C. The gene lies in close proximity to another family
	member (TRPV1) gene on chromosome 17, and the two encoded proteins are thought to
	associate with each other to form heteromeric channels (1, 2).
Gene ID:	497948
NCBI Accession:	NP_001020928
UniProt:	Q4QYD9
Application Details	
Application Notes:	• WB (1:1000)
	• IHC (1:1000)
	• ICC/IF (1:100)
	optimal dilutions for assays should be determined by the user.
Comment:	1 μ g/ml of ABIN2485362 was sufficient for detection of TrpV3 in 10 μ g of COS-1 cell lysate
	transiently transfected with TrpV3 by colorimetric immunoblot analysis using Goat anti-mouse
	IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C







Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone N15/4 (ABIN2485362). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody (ABIN2485362) at 1:100 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) TrpV3 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Human Cell lysates showing detection of TrpV3 protein using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-4. Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-TrpV3 Monoclonal Antibody, Clone S15-4. Tissue: hippocampus. Species: Human. Fixation: Bouin's Fixative and paraffin-embedded. Primary Antibody: Mouse Anti-TrpV3 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT.

Please check the product details page for more images. Overall 4 images are available for ABIN2485362.