antibodies -online.com





Datasheet for ABIN2485377

anti-TRPC5 antibody (AA 827-845) (APC)

4 Images



Go to Product page

_						
0	V	е.	r_{V}	/	6	W


Quantity:	100 μg	
Target:	TRPC5	
Binding Specificity:	AA 827-845	
Reactivity:	Human	
Host:	Mouse	
Clonality:	Monoclonal	
Conjugate:	This TRPC5 antibody is conjugated to APC	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Antibody Array (AA)	


Product Details


Immunogen:	Synthetic peptide amino acids 827-845 of human TrpC5 (also known as short transient receptor potential channel 5, and Htrp5)
Clone:	N67-15 (Formerly S67-15)
Isotype:	lgG2b
Specificity:	Detects ~110 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

Target Details

900 2 0 0		
Target:	TRPC5	
Alternative Name:	TRPC5 (TRPC5 Products)	
Background:	Transient receptor potential cation channel, subfamily C, member 5, also known as TRPC5, is a subtype of the TRPC family of mammalian transient receptor potential ion channels. Homomultimeric TRPC5 and hetero-multimeric TRPC5-TRPC1 channels are activated by extracellula reduced thioredoxin (1). This activation probably plays a role in rheumatoid arthritis. It has also been recently found to be involved in the action on anaesthetics such as chloroform, halothane and propofol (2).	
Gene ID:	7224	
NCBI Accession:	NP_036603	
UniProt:	Q9UL62	
Application Details		
Application Notes:	 WB (1:1000) IHC (1:1000) ICC/IF (1:100) optimal dilutions for assays should be determined by the user. 	
Comment:	1 μg/ml of ABIN2485377 was sufficient for detection of TrpC5 in 20 μg of rat brain lysate by colorimetric immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Conjugated antibodies should be stored at 4°C	







Western Blotting

Image 1. Western Blot analysis of Mouse brain showing detection of 110 kDa TrpC5 protein using Mouse Anti-TrpC5 Monoclonal Antibody, Clone N67/15 (ABIN2485377). Lane 1: Molecular Weight Ladder (MW). Lane 2: Mouse Brain. Load: 15 µg. Block: 5 % Skim Milk powder in TBST. Primary Antibody: Mouse Anti-TrpC5 Monoclonal Antibody (ABIN2485377) at 1:1000 for Overnight at 4C. Secondary Antibody: Goat anti-mouse IgG:HRP at 1:7000 for 1 hour at RT with shaking. Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT. Predicted/Observed Size: 110 kDa.

Western Blotting

Image 2. Western Blot analysis of Rat Brain Membrane showing detection of ~110 kDa TrpC5 protein using Mouse Anti-TrpC5 Monoclonal Antibody, Clone S67-15 . Lane 1: Molecular Weight Ladder (MW). Lane 2: Rat Brain Membrane cell lysate. Load: 20 μg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-TrpC5 Monoclonal Antibody at 1:1000 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:100 for 60 min at RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~110 kDa. Other Band(s): 160kDa, 60kDa.

Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-TrpC5 Monoclonal Antibody, Clone S67-15. Tissue: Brain Slice. Species: Mouse. Fixation: 10% Formalin Solution for 12-24 hours at RT. Primary Antibody: Mouse Anti-TrpC5 Monoclonal Antibody at 1:1000 for 1 hour at RT. Secondary Antibody: HRP/DAB Detection System: Biotinylated Goat Anti-Mouse, Streptavidin Peroxidase, DAB Chromogen (brown) for 30 minutes at RT. Counterstain: Mayer

Hematoxylin (purple/blue) nuclear stain at 250-500 μ l for 5 minutes at RT. Localization: Nuclear staining. Magnification: 10X.

Please check the product details page for more images. Overall 4 images are available for ABIN2485377.