# antibodies .- online.com







## anti-SHANK3 antibody (AA 840-857) (HRP)



**Images** 



| $\sim$ |          |
|--------|----------|
| ( )\/△ | rview    |
| $\cup$ | 1 410 44 |

| Quantity:            | 100 μg   |
|----------------------|--|
| Target:              | SHANK3   |
| Binding Specificity: | AA 840-857   |
| Reactivity:          | Rat  |
| Host:                | Mouse  |
| Clonality:           | Monoclonal   |
| Conjugate:           | This SHANK3 antibody is conjugated to HRP  |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC), Immunoprecipitation (IP), Immunofluorescence (IF), Immunocytochemistry (ICC), Antibody Array (AA) |

## **Product Details**

| Immunogen:        | Synthetic peptide amino acids 840-857 of rat Shank3             |
|-------------------|---|
| Clone:            | S69   |
| Isotype:          | lgG2b   |
| Specificity:      | Detects ~190 kDa. No cross-reactivity against Shank1 or Shank2. |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Protein G Purified  |

## **Target Details**

Target: SHANK3


## **Target Details**

| - Target Details    |  |
|---------------------|--|
| Alternative Name:   | SHANK3 (SHANK3 Products)   |
| Background:         | Shank proteins make up a family of scaffold proteins identified through their interaction with a           |
|                     | variety of membrane and cytoplasmic proteins (1). Shank proteins at postsynaptic sites of                  |
|                     | excitatory synapses play roles in signal transmission into the postsynaptic neuron. Shank                  |
|                     | proteins are also crucial in receptor tyrosine kinase signaling, specifically, Shank3 can mediate          |
|                     | Erk-MAPK and P13K signaling which is crucial for tubule formation (2). Shank3 is also one of               |
|                     | the latest genes to be associated with autism. A mutation of a single copy of Shank3 on                    |
|                     | chromosome 22q13 can result in language and/or social communication disorders (3).                         |
| Gene ID:            | 59312  |
| NCBI Accession:     | NP_067708  |
| UniProt:            | Q9JLU4   |
| Pathways:           | Synaptic Membrane, Tube Formation, Regulation of long-term Neuronal Synaptic Plasticity                    |
| Application Details |  |
| Application Notes:  | • WB (1:1000)  |
|                     | • IHC (1:100)  |
|                     | <ul> <li>ICC/IF (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul> |
|                     | Optimal dilutions for assays should be determined by the user.   |
| Comment:            | 1 μg/ml of ABIN2485414 was sufficient for detection of Shank3 in 10 μg COS cell lysate                     |
|                     | transiently transfected with Shank3 by colorimetric immunoblot analysis using goat anti-mouse              |
|                     | IgG:HRP as the secondary antibody.   |
| Restrictions:       | For Research Use only  |
| Handling            |  |
| Format:             | Liquid   |
| Concentration:      | 1 mg/mL  |
| Buffer:             | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated                  |
| Preservative:       | Sodium azide   |
| Precaution of Use:  | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which                              |
|                     | should be handled by trained staff only.   |
| Storage:            | 4 °C   |
|                     |  |

Storage Comment:


Conjugated antibodies should be stored at 4°C

## **Images**




#### **Immunocytochemistry**

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SHANK3 Monoclonal Antibody, Clone S69-46 (ABIN2485414). Tissue: HaCaT cells. Species: Human. Fixation: Cold 100 % methanol for 10 minutes at -20 °C. Primary Antibody: Mouse Anti-SHANK3 Monoclonal Antibody (ABIN2485414) at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Borderline positive.



### Immunofluorescence (fixed cells)

Image 2. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-SHANK3 Monoclonal Antibody, Clone S69-46. Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol for 10 minutes at -20°C. Primary Antibody: Mouse Anti-SHANK3 Monoclonal Antibody at 1:100 for 1 hour at RT. Secondary Antibody: FITC Goat Anti-Mouse (green) at 1:50 for 1 hour at RT. Localization: Borderline positive.



#### **Western Blotting**

**Image 3.** Western Blot analysis of Rat brain membrane lysate showing detection of SHANK3 protein using Mouse Anti-SHANK3 Monoclonal Antibody, Clone S69-46. Load: 15 μg. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Mouse Anti-SHANK3 Monoclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Sheep Anti-Mouse IgG: HRP for 1 hour at RT.

Please check the product details page for more images. Overall 5 images are available for ABIN2485414.