# antibodies - online.com







## anti-CLCN3 antibody (AA 98-115) (Atto 594)





$\sim$	
( )\/\p	rview
$\circ$	

Quantity:	100 μg
Target:	CLCN3
Binding Specificity:	AA 98-115
Reactivity:	Rat
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This CLCN3 antibody is conjugated to Atto 594
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)


#### **Product Details**


Immunogen:	Synthetic peptide amino acids 98-115 (cytoplasmic N-terminus) of rat Clcn3
Clone:	S258-5
Isotype:	IgG1
Specificity:	Detects ~90 kDa. Does not cross-react with Clcn4 or Clcn5 (based on KO validation results).
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

## **Target Details**

## **Target Details**

rarget Details		
Alternative Name:	CICN3 (CLCN3 Products)	
Background:	Clcn3 mediates the exchange of chloride ions against protons, and functions as an antiporter and contributes to the acidification of the endosome and synaptic vesicle lumen, and may thereby affect vesicle trafficking and exocytosis. It may play an important role in neuronal cell function through regulation of membrane excitability by protein kinase C. It could also help neuronal cells to establish short-term memory.	
Gene ID:	84360	
NCBI Accession:	NP_445815	
UniProt:	P51792	
Application Details		
Application Notes:	<ul> <li>WB (1:1000)</li> <li>IHC (1:1000)</li> <li>ICC/IF (1:100)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>	
Comment:	1 $\mu$ g/ml of ABIN2485473 was sufficient for detection of Clcn3 in 20 $\mu$ g of rat brain membrane lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 mg/mL	
Buffer:	PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated	
Preservative:	Sodium azide	
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.	
Storage:	4 °C	
Storage Comment:	Conjugated antibodies should be stored at 4°C	






#### **Immunocytochemistry**

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Clcn3 Monoclonal Antibody, Clone S258-5 (ABIN2485473). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Clcn3 Monoclonal Antibody (ABIN2485473) at 1:50 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Clcn3 Antibody (D) Composite.

#### Immunofluorescence (fixed cells)

Immunocytochemistry/Immunofluorescence 2. **Image** analysis using Mouse Anti-Clcn3 Monoclonal Antibody, Clone S258-5. Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Clcn3 Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Membrane, Endosome, Endosome membrane, Vesicle, Secretory Vesicle Membrane. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Clcn3 Antibody (D) Composite.



### **Western Blotting**

Image 3. Western Blot analysis of Rat Brain Membrane showing detection of ~90 kDa Clcn3 protein using Mouse Anti-Clcn3 Monoclonal Antibody, Clone S258-5 . Lane 1: Molecular Weight Ladder. Lane 2: Rat Brain Membrane. Load: 15 μg. Block: 2% BSA and 2% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-Clcn3 Monoclonal Antibody at 1:200 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: ECL solution for 6 min in RT. Predicted/Observed Size: ~90 kDa.