

Datasheet for ABIN2485746  
**anti-Versican antibody (AA 362-585) (FITC)**[Go to Product page](#)

## 3 Images

## Overview

Quantity:	100 µg
Target:	Versican (Vcan)
Binding Specificity:	AA 362-585
Reactivity:	Mouse
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Versican antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunofluorescence (IF)

## Product Details

Immunogen:	Fusion protein amino acids 362-585 (glycosaminoglycan alpha domain) of mouse Versican core protein
Clone:	S351-23
Isotype:	IgG1
Specificity:	Detects >350 kDa.
Cross-Reactivity:	Human, Mouse, Rat
Purification:	Protein G Purified

## Target Details

Target:	Versican (Vcan)
---------	-----------------

## Target Details

Alternative Name: Versican ([Vcan Products](#))

Background: Versican (chondroitin sulfate proteoglycan 2) is a large extracellular matrix proteoglycan involved in cell growth and differentiation. Important as a structural molecule, versican creates loose and hydrated matrices during key events in development and disease. The protein contains hyaluronic acid and glycosaminoglycan-binding domains, epidermal growth factor-like repeats, a Lectinlike sequence and a complement regulatory protein-like domain. Splice variants differ greatly in length and degree of modification by glycoaminoglycan chains. Accumulation around smooth muscle cells in lesions of athero-sclerosis suggests a role for versican in atherogenesis. Versican, differentially expressed in human melanoma, plays a role in tumor development and may be a reliable marker for clinical diagnosis. The organization of HA- and versican-rich pericellular matrices may facilitate migration and mitosis by diminishing cell surface adhesivity and affecting cell shape through steric exclusion and the viscous properties of HA proteoglycan gels.

Gene ID: 13003

UniProt: [Q62059](#)

Pathways: [Glycosaminoglycan Metabolic Process](#)

## Application Details

Application Notes:

- WB (1:1000)
- ICC/IF (1:100)
- optimal dilutions for assays should be determined by the user.

Comment: 1 µg/ml of ABIN2485746 was sufficient for detection of Versican in 20 µg of mouse brain membrane lysate and assayed by colorimetric immunoblot analysis using goat anti-mouse IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 mg/mL


Buffer: PBS pH 7.4, 50 % glycerol, 0.1 % sodium azide, Storage buffer may change when conjugated

Preservative: Sodium azide

Handling


Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C

Images




Immunocytochemistry

**Image 1.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Versican Monoclonal Antibody, Clone S351-23 (ABIN2485746). Tissue: Neuroblastoma cells (SH-SY5Y). Species: Human. Fixation: 4 % PFA for 15 min. Primary Antibody: Mouse Anti-Versican Monoclonal Antibody (ABIN2485746) at 1:200 for overnight at 4 °C with slow rocking. Secondary Antibody: AlexaFluor 488 at 1:1000 for 1 hour at RT. Counterstain: Phalloidin-iFluor 647 (red) F-Actin stain, Hoechst (blue) nuclear stain at 1:800, 1.6 mM for 20 min at RT. (A) Hoechst (blue) nuclear stain. (B) Phalloidin-iFluor 647 (red) F-Actin stain. (C) Versican Antibody (D) Composite.



Immunofluorescence (fixed cells)

**Image 2.** Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-Versican Monoclonal Antibody, Clone S351-23 . Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4% Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-Versican Monoclonal Antibody at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain; DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60min RT, 5min RT. Localization: Cytoplasm, Membrane, Extracellular Space, Extracellular Matrix. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) Versican Antibody (D) Composite.



**Western Blotting**

**Image 3.** Western Blot analysis of Rat Brain Membrane and brain showing detection of 350kDa Versican protein using Mouse Anti-Versican Monoclonal Antibody, Clone S351-23 . Lane 1: Molecular Weight Ladder. Lane 2: Rat Brain Membrane and brain. Load: 15 µg. Block: 5% Skim Milk in 1X TBST. Primary Antibody: Mouse Anti-Versican Monoclonal Antibody at 1:200 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:1000 for 1 hour RT. Color Development: KPL TMB. Predicted/Observed Size: 350kDa. Other Band(s): Multiple other bands.