

Datasheet for ABIN2485816
anti-NPAS4 antibody (AA 597-802) (APC)[Go to Product page](#)

2 Images

Overview

| | |
|----------------------|--|
| Quantity: | 100 µg |
| Target: | NPAS4 |
| Binding Specificity: | AA 597-802 |
| Reactivity: | Rat |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This NPAS4 antibody is conjugated to APC |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC) |

Product Details

| | |
|-------------------|--|
| Immunogen: | Fusion protein amino acids 597-802 (C-terminus) of rat Npas4. |
| Clone: | S408-79 |
| Isotype: | IgG1 |
| Specificity: | Highly expressed in the brain, with lower expression found in endocrine tissue., Detects ~90 kDa. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Protein G Purified |

Target Details


| | |
|-------------------|--|
| Target: | NPAS4 |
| Alternative Name: | NPAS4 (NPAS4 Products) |
| Background: | <p>NPAS4 belongs to the Per-Arnt-Sim family of neuronal specific transcription factors, all involved in the development and maintenance of learning and memory pathways (1). NPAS4 is a transcription factor that regulates the formation and maintenance of inhibitory synapses, and can activate the CNS midline enhancer element and the expression of the drebrin gene.</p> <p>Deregulation has been to be associated with developmental disorders such as schizophrenia and autism (2).</p> |
| Gene ID: | 266734 |
| NCBI Accession: | NP_705890 |
| UniProt: | Q8CJH6 |

Application Details

| | |
|--------------------|--|
| Application Notes: | <ul style="list-style-type: none">• WB (1:1000)• ICC/IF(1:100)• optimal dilutions for assays should be determined by the user. |
| Comment: | A 1:100 dilution of ABIN2485816 was sufficient for detection of NPAS4 in 20 µg of mouse brain lysate by ECL immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody. |
| Restrictions: | For Research Use only |


Handling

| | |
|--------------------|--|
| Format: | Liquid |
| Concentration: | 1 mg/mL |
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.1 % sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Conjugated antibodies should be stored at 4°C |



Immunocytochemistry

Image 1. Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-NPAS4 Monoclonal Antibody, Clone S408-79 (ABIN2485816). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4 % Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-NPAS4 Monoclonal Antibody (ABIN2485816) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain, DAPI (blue) nuclear stain at 1:1000, 1:5000 for 60 min RT, 5 min RT. Localization: Cytoplasm. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) NPAS4 Antibody (D) Composite.



Western Blotting

Image 2. Western Blot analysis of Rat Brain showing detection of ~90 kDa NPAS4 protein using Mouse Anti-NPAS4 Monoclonal Antibody, Clone S408-79 . Lane 1: MW Ladder. Lane 2: Rat Brain. Load: 20 µg. Block: 2% GE Healthcare Blocker for 1 hour at RT. Primary Antibody: Mouse Anti-NPAS4 Monoclonal Antibody at 1:1000 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:200 for 1 hour at RT. Color Development: ECL solution for 6 min at RT. Predicted/Observed Size: ~90 kDa. Other Band(s): ~60, 45, 40, 38, 25, 20 kDa.