antibodies -online.com





anti-NPAS4 antibody (AA 597-802) (FITC)

2 Images



Go to Product page

Overview


| Quantity: | 100 μg |
|----------------------|---|
| Target: | NPAS4 |
| Binding Specificity: | AA 597-802 |
| Reactivity: | Rat |
| Host: | Mouse |
| Clonality: | Monoclonal |
| Conjugate: | This NPAS4 antibody is conjugated to FITC |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunocytochemistry (ICC) |


Product Details

| Immunogen: | Fusion protein amino acids 597-802 (C-terminus) of rat Npas4. |
|-------------------|--|
| Clone: | S408-79 |
| Isotype: | lgG1 |
| Specificity: | Highly expressed in the brain, with lower expression found in endocrine tissue.,Detects ~90 kDa. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Protein G Purified |

Target Details

| Target: | NPAS4 | |
|---------------------|--|--|
| Alternative Name: | NPAS4 (NPAS4 Products) | |
| Background: | NPAS4 belongs to the Per-Arnt-Sim family of neuronal specific transciption factors, all involved in the development and maintenance of learning and memory pathways (1). NPAS4 is a transcription factor that regulates the formation and maintenance of inhibitory synapses, and can activate the CNS midline enhancer element and the expression of the drebrin gene. Deregulation has been to be associated with developmental disorders such as schizophrenia and autism (2). | |
| Gene ID: | 266734 | |
| NCBI Accession: | NP_705890 | |
| UniProt: | Q8CJH6 | |
| Application Details | | |
| Application Notes: | WB (1:1000) ICC/IF(1:100) optimal dilutions for assays should be determined by the user. | |
| Comment: | A 1:100 dilution of ABIN2485818 was sufficient for detection of NPAS4 in 20 µg of mouse brain lysate by ECL immunoblot analysis using Goat anti-mouse IgG:HRP as the secondary antibody. | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 mg/mL | |
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.1 % sodium azide, Storage buffer may change when conjugated | |
| Preservative: | Sodium azide | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. | |
| Storage: | 4 °C | |
| Storage Comment: | Conjugated antibodies should be stored at 4°C | |





Immunocytochemistry

Immunocytochemistry/Immunofluorescence analysis using Mouse Anti-NPAS4 Monoclonal Antibody, Clone S408-79 (ABIN2485818). Tissue: Neuroblastoma cell line (SK-N-BE). Species: Human. Fixation: 4 % Formaldehyde for 15 min at RT. Primary Antibody: Mouse Anti-NPAS4 Monoclonal Antibody (ABIN2485818) at 1:100 for 60 min at RT. Secondary Antibody: Goat Anti-Mouse ATTO 488 at 1:100 for 60 min at RT. Counterstain: Phalloidin Texas Red F-Actin stain, DAPI (blue) nuclear stain at 1:1000, 1:5000 for 5 min RT. Localization: Cytoplasm. Magnification: 60X. (A) DAPI (blue) nuclear stain (B) Phalloidin Texas Red F-Actin stain (C) NPAS4 Antibody (D) Composite.

Western Blotting

Image 2. Western Blot analysis of Rat Brain showing detection of ~90 kDa NPAS4 protein using Mouse Anti-NPAS4 Monoclonal Antibody, Clone S408-79 . Lane 1: MW Ladder. Lane 2: Rat Brain. Load: 20 μg. Block: 2% GE Healthcare Blocker for 1 hour at RT. Primary Antibody: Mouse Anti-NPAS4 Monoclonal Antibody at 1:1000 for 16 hours at 4°C. Secondary Antibody: Goat Anti-Mouse IgG: HRP at 1:200 for 1 hour at RT. Color Development: ECL solution for 6 min at RT. Predicted/Observed Size: ~90 kDa. Other Band(s): ~60, 45, 40, 38, 25, 20 kDa.