
antibodies -online.com





anti-Nitrotyrosine antibody (Atto 594)



Images



Overview

Quantity:	100 μg
Target:	Nitrotyrosine
Reactivity:	Please inquire
Host:	Mouse
Clonality:	Monoclonal
Conjugate:	This Nitrotyrosine antibody is conjugated to Atto 594
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC), Immunoprecipitation (IP), Flow Cytometry (FACS), Immunocytochemistry (ICC), Immunofluorescence (IF), Antibody Array (AA)

Product Details

Immunogen:	3-(4-hydroxy-3-nitrophenylacetamido) propionic acid-bovine serum albumin
Clone:	39B6
Isotype:	lgG2a
Specificity:	Recognizes 3-nitrotyrosine moieties. No detectable cross-reactivity with non-nitrated tyrosine. Not species specific.
Purification:	Protein G Purified

Target Details

Target:	Nitrotyrosine
Abstract:	Nitrotyrosine Products

Target Details

Target Type:	Chemical
Background:	Protein tyrosine nitration results in a post-translational modification that is increasingly
	receiving attention as an important component of nitric oxide signaling (2). While multiple
	nonenzymatic mechanisms are known to be capable of producing nitrated tyrosine residues,
	most tyrosine nitration events involve catalysis by metalloproteins such as myeloperoxidase
	eosino-philperoxidase (3), myoglobin, the cytochrome P-450s, superoxide dismutase and
	prostacyclin synthase. Nitrotyrosine may also serve as a biomarker for the effects of reactive
	nitrogen oxides, based on tyrosine residues becoming nitrated in proteins at sites of
	inflammation induced tissue injury (1). The presence of nitro tyrosine-containing proteins
	therefore has shown high correlation to disease states such as atherosclerosis, Alzheimer's
	disease, Parkinson's disease and amyotrophic lateral sclerosis (4).
Application Details	
Application Notes:	• WB (1:1400)
	• IHC (1:100)
	 optimal dilutions for assays should be determined by the user.
Comment:	0.7 μg/ml of ABIN2486185 was sufficient for detection of 5 μg SIN-1 treated BSA by Western
	Blot analysis using Goat anti-mouse IgG:HRP as the secondary antibody.
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which
	should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C








Image 1. Western Blot analysis of Human Recombinant Protein showing detection of Multiple Bands Nitrotyrosine protein using Mouse Anti-Nitrotyrosine Monoclonal Antibody, Clone 39B6 (ABIN2486185). Lane 1: MW Ladder. Lane 2: hASYN Monomer (3.84 μg). Lane 3: Nitrosylated hASYN (3.84 μg).. Block: 5 % Skim Milk Powder in TBST. Primary Antibody: Mouse Anti-Nitrotyrosine Monoclonal Antibody (ABIN2486185) at 1:1000 for 2 hours at RT with shaking . Secondary Antibody: Goat anti-mouse IgG:HRP at 1:4000 for 2 hour at RT with shaking . Color Development: Chemiluminescent for HRP (Moss) for 5 min in RT. Predicted/Observed Size: Multiple Bands.



Immunohistochemistry

Image 2. Immunohistochemistry analysis using Mouse Anti-Nitrotyrosine Monoclonal Antibody, Clone 39B6. Tissue: inflamed colon. Species: Mouse. Fixation: Formalin. Primary Antibody: Mouse Anti-Nitrotyrosine Monoclonal Antibody at 1:1000000 for 12 hours at 4°C. Secondary Antibody: Biotin Goat Anti-Mouse at 1:2000 for 1 hour at RT. Counterstain: Mayer Hematoxylin (purple/blue) nuclear stain at 200 µl for 2 minutes at RT. Magnification: 40x.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Mouse Anti-Nitrotyrosine Monoclonal Antibody, Clone 39B6. Tissue: liver tissue. Species: Rat. Primary Antibody: Mouse Anti-Nitrotyrosine Monoclonal Antibody at 1:1000. Secondary Antibody: FITC Goat Anti-Mouse (green).

Please check the product details page for more images. Overall 7 images are available for ABIN2486185.