antibodies -online.com





anti-SLC12A3 antibody (AA 74-95) (APC)






| Overview | | |
|----------------------|--|--|
| Quantity: | 100 μg | |
| Target: | SLC12A3 | |
| Binding Specificity: | AA 74-95 | |
| Reactivity: | Rat | |
| Host: | Rabbit | |
| Clonality: | Polyclonal | |
| Conjugate: | This SLC12A3 antibody is conjugated to APC | |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF), Immunoelectron Microscopy (IEM) | |
| Product Details | | |
| Immunogen: | Produced against a synthetic peptide mapping to a segment of rat NCC (amino acids 74-95), N-terminal | |
| Specificity: | Detects ~160 kDa. | |
| Cross-Reactivity: | Dog, Human, Mouse, Rat | |
| Purification: | Affinity Purified | |
| Target Details | | |
| Target: | SLC12A3 | |
| Alternative Name: | NCC (SLC12A3 Products) | |


Target Details

| Background: | NCC, a thiazide-sensitive NaCl co-transporter, is found on the apical membrane of the distal convoluted tubule, where it is the principal mediator of Na+ and Cl- reabsorption in this | | |
|---------------------|--|--|--|
| | segment of the nephron. It is activated by phosphorylation, and has been implicated in renal | | |
| | NaCl and K+ homeostasis (1). Regulation of NCC expression and phosphorylation by dietary | | |
| | NACI restriction appears to involve SGK1(1). In experiments with angiotensin II-infused mice, | | |
| | increased sensitivity to Ang II may involved over-activity of NCC (2). Therefore, NCC is the | | |
| | target of thiazide diuretics used in the treatment of hypertension (1). Molecular experiments | | |
| | have also shown that NCC has been detected in the lens cortex, core and fiber cells of a rat (3) | | |
| Gene ID: | 54300, 6559 | | |
| NCBI Accession: | NP_062218, NP_000330 | | |
| UniProt: | P55018, P55017 | | |
| Application Details | | | |
| Application Notes: | • WB (1:1000) | | |
| | • IHC (1:200) | | |
| | optimal dilutions for assays should be determined by the user. | | |
| Comment: | 1 μg/ml of ABIN2486291 was sufficient for detection of NCC3 in 10 μg of rat kidney tissue | | |
| | lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary | | |
| | antibody. | | |
| Restrictions: | For Research Use only | | |
| Handling | | | |
| Format: | Liquid | | |
| Concentration: | 1 mg/mL | | |
| Buffer: | PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated | | |
| Preservative: | Sodium azide | | |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which | | |
| | should be handled by trained staff only. | | |
| Storage: | 4 °C | | |
| Storage Comment: | Conjugated antibodies should be stored at 4°C | | |
| | | | |




Immunohistochemistry

Image 1. Immunohistochemistry analysis using Rabbit Anti-NCC Polyclonal Antibody (ABIN2486291). Tissue: kidney tissue. Species: Mouse. Fixation: paraformaldehyde-fixed paraffin-embedded. Primary Antibody: Rabbit Anti-NCC Polyclonal Antibody (ABIN2486291) at 1:1000 for Overnight at 4 °C. Secondary Antibody: Biotinylated Goat Anti-Rabbit IgG at 1:500 for 30 min at RT. Courtesy of: Brigitte Rogge.



Western Blotting

Image 2. Western blot analysis of Rat tissue lysates showing detection of NCC protein using Rabbit Anti-NCC Polyclonal Antibody . Primary Antibody: Rabbit Anti-NCC Polyclonal Antibody at 1:1000.



Immunohistochemistry

Image 3. Immunohistochemistry analysis using Rabbit Anti-NCC Polyclonal Antibody . Tissue: kidney tissue. Species: Rat. Primary Antibody: Rabbit Anti-NCC Polyclonal Antibody at 1:200. Secondary Antibody: FITC Goat Anti-Rabbit (green).

Please check the product details page for more images. Overall 4 images are available for ABIN2486291.