

Datasheet for ABIN2486325  
**anti-SLC12A1 antibody (AA 33-55) (APC)**[Go to Product page](#)

## 2 Images

## Overview

|                      |   |
|----------------------|---|
| Quantity:            | 100 µg  |
| Target:              | SLC12A1   |
| Binding Specificity: | AA 33-55  |
| Reactivity:          | Rat   |
| Host:                | Rabbit  |
| Clonality:           | Polyclonal  |
| Conjugate:           | This SLC12A1 antibody is conjugated to APC        |
| Application:         | Western Blotting (WB), Immunohistochemistry (IHC) |

## Product Details

|                   |  |
|-------------------|--|
| Immunogen:        | Produced against a synthetic peptide mapping to a segment (amino acids 33-55) of the N-terminal tail of rat NKCC2 (antibody designation L224). |
| Specificity:      | Detects ~160 kDa.  |
| Cross-Reactivity: | Human, Mouse, Rat  |
| Purification:     | Protein A Purified   |

## Target Details

|                   |   |
|-------------------|---|
| Target:           | SLC12A1   |
| Alternative Name: | NKCC2 ( <a href="#">SLC12A1 Products</a> )  |
| Background:       | NKCC2, a kidney-specific member of the cation chloride co-transporter family SLC12, facilitates |

## Target Details

transport of sodium, potassium, and chloride ions from the lumen of the renal thick ascending limb into the cell (1, 2). Transport of sodium dilutes the luminal fluid, decreasing its osmolality creating an osmotic driving force for water reabsorption in the connecting tubule and cortical collecting duct under the influence of the hormone vasopressin. NKCC2 is found on the apical surface of the thick ascending limb of the loop of Henle and is blocked by loop diuretics such as furosemide (1).

Gene ID: 25065

NCBI Accession: [NP\\_001257546](#)

UniProt: [Q5PQV1](#)

## Application Details

Application Notes:

- WB (1:1000)
- IHC (1:25)
- optimal dilutions for assays should be determined by the user.

Comment: 1 µg/ml of ABIN2486325 was sufficient for detection of NKCC2 in 10 µg of rat kidney tissue lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 mg/mL


Buffer: PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated

Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.


Storage: 4 °C

Storage Comment: Conjugated antibodies should be stored at 4°C



**Immunohistochemistry**

**Image 1.** Immunohistochemistry analysis using Rabbit Anti-NKCC2 Polyclonal Antibody (ABIN2486325). Tissue: Kidney. Species: Rat. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-NKCC2 Polyclonal Antibody (ABIN2486325) at 1:25 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit IgG: Alexa Fluor 488. Counterstain: Actin-binding Phalloidin-Alexa Fluor 633, DAPI (blue) nuclear stain. Magnification: 63X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Alex Fluor 633 F-Actin stain. (C) NKCC2 Antibody (D) Composite



**Western Blotting**

**Image 2.** Western blot analysis of Rat kidney tissue lysates showing detection of NKCC2 protein using Rabbit Anti-NKCC2 Polyclonal Antibody . Primary Antibody: Rabbit Anti-NKCC2 Polyclonal Antibody at 1:1000.