

## Datasheet for ABIN2486394

# anti-Aquaporin 4 antibody (C-Term) (Biotin)

2 Images




Go to Product page

_			
	Ve.	rv	iew

Quantity:	100 μg	
Target:	Aquaporin 4 (AQP4)	
Binding Specificity:	C-Term	
Reactivity:	Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Aquaporin 4 antibody is conjugated to Biotin	
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunocytochemistry (ICC), Immunofluorescence (IF)	
Product Details		
Product Details Immunogen:	Produced against the C-terminal peptide (Sequence N-CTKGSYMEVEDNRSQVETED) of rat aquaporin 4	
Immunogen:	aquaporin 4	
Immunogen: Specificity:	aquaporin 4  Detects ~35 kDa. May detect larger glycosylated bands ~50 kDa.	
Immunogen:  Specificity:  Cross-Reactivity:	aquaporin 4  Detects ~35 kDa. May detect larger glycosylated bands ~50 kDa.  Human, Mouse, Rat	
Immunogen:  Specificity:  Cross-Reactivity:  Purification:	aquaporin 4  Detects ~35 kDa. May detect larger glycosylated bands ~50 kDa.  Human, Mouse, Rat	


## **Target Details**

-			
Background:	Aquaporins selectively conduct water molecules in and out of the cell, while preventing the		
	passage of ions and other solutes. Known as water channels, they are integral membrane pore		
	proteins (1, 2). Aquaporin 4 is constitutively expressed in the basolateral cell membrane of		
	principal collecting duct cells in the kidney and provide a pathway for water to exit these cells		
	(3). It is also expressed in astrocytes and is up regulated by direct insult to the CNS (4).		
	Aquaporin 4 has been proposed as the primary autoimmune target in neuromyelitis optica (5,		
	6).		
Gene ID:	25293		
NCBI Accession:	NP_036957		
UniProt:	P47863		
Pathways:	Sensory Perception of Sound		
Application Details			
Application Notes:	• WB (1:2000)		
	• IHC (1:200)		
	ICC/IF (1:400)      antime! dilutions for account should be determined by the user.		
	optimal dilutions for assays should be determined by the user.		
Comment:	0.5 μg/ml of ABIN2486394 was sufficient for detection of aquaporin 4 in 10 μg of rat kidney		
	tissue lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the		
	secondary antibody.		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 mg/mL		
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated		
Preservative:	Sodium azide		
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which		
	should be handled by trained staff only.		
Storage:	4 °C		
Storage Comment:	Conjugated antibodies should be stored at 4°C		
	oonjugated antibodies should be stored at 4 o		



#### Immunohistochemistry

**Image 1.** Immunohistochemistry analysis using Rabbit Anti-Aquaporin 4 Polyclonal Antibody . Tissue: kidney tissue. Species: Rat. Primary Antibody: Rabbit Anti-Aquaporin 4 Polyclonal Antibody at 1:200. Secondary Antibody: FITC Goat Anti-Rabbit (green).



### **Western Blotting**

**Image 2.** Western blot analysis of Rat kidney inner medullary homogenates showing detection of Aquaporin 4 protein using Rabbit Anti-Aquaporin 4 Polyclonal Antibody. Primary Antibody: Rabbit Anti-Aquaporin 4 Polyclonal Antibody at 1:2000.