antibodies .- online.com







anti-SCNN1A antibody (AA 617-638) (Atto 488)



Images



-						
O	V	e	rv	1	е	W

Quantity:	100 μg		
Target:	SCNN1A		
Binding Specificity:	AA 617-638		
Reactivity:	Rat		
Host:	Rabbit		
Clonality:	Polyclonal		
Conjugate:	This SCNN1A antibody is conjugated to Atto 488		
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),		
	Immunoprecipitation (IP), Immunocytochemistry (ICC)		
Product Details			
Immunogen:	Produced against the C-terminal tail (amino acids 617-638) of rat beta ENaC (antibody		
	designation 3755-2)		
Specificity:	Detects ~87 kDa.		
Cross-Reactivity:	Hamster, Human, Mouse, Rat, Xenopus laevis		
Purification:	Protein A Purified		
Target Details			
Target:	SCNN1A		
Alternative Name:	ENaC (SCNN1A Products)		


Target Details

Background:	The Epithelial Sodium Channel (ENaC) is a membrane ion channel permeable to Na+ ions. It is				
	located in the apical plasma membrane of epithelia in the kidneys, lung, colon, and other tissues				
	where it plays a role in trans epithelial Na+-ion transport (1). Specifically Na+ transport via ENaC				
	occurs across many epithelial surfaces, and plays a key role in regulating salt and water				
	absorption (2). ENaCs are composed of three structurally related subunits that form a				
	tetrameric channel, α , β , and γ . The expression of its alpha and beta subunits is enhanced as				
	keratinocytes differentiate (3, 4). The beta and gamma-ENaC subunits are essential for edema				
	fluid to exert its maximal effect on net fluid absorption by distal lung epithelia(5). And it has				
	been concluded that the subunits are differentially expressed in the retina of mice with ocular				
	hypertension, therefore the up-regulation of alpha-ENaC proteins could serve as a protection				
	mechanism against elevated intraocular pressure (6).				
Gene ID:	24767				
NCBI Accession:	NP_036780				
UniProt:	P37090				
Application Details					
Application Notes:	• WB (1:1000)				
	• IHC (1:100)				
	optimal dilutions for assays should be determined by the user.				
Comment:	1 μg/ml of ABIN2486401 was sufficient for detection of beta-ENaC in 20 μg of rat kidney tissue				
	lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary				
	antibody.				
Restrictions:	For Research Use only				
Handling					
Format:	Liquid				
Concentration:	1 mg/mL				
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated				
Preservative:	Sodium azide				
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which				
	should be handled by trained staff only.				

Storage: 4 °C


Storage Comment: Conjugated antibodies should be stored at 4°C

Images




Immunohistochemistry

Image 1. Immunohistochemistry analysis using Rabbit Anti-ENaC Polyclonal Antibody . Tissue: kidney tissue. Species: Rat. Primary Antibody: Rabbit Anti-ENaC Polyclonal Antibody at 1:100. Secondary Antibody: FITC Goat Anti-Rabbit (green).



Western Blotting

Image 2. Western blot analysis of Rat kidney tissue lysates showing detection of ENaC protein using Rabbit Anti-ENaC Polyclonal Antibody . Primary Antibody: Rabbit Anti-ENaC Polyclonal Antibody at 1:1000.



Western Blotting

Image 3. Western blot analysis of Mouse mpkCCD cell lysates showing detection of ENaC protein using Rabbit Anti-ENaC Polyclonal Antibody . Primary Antibody: Rabbit Anti-ENaC Polyclonal Antibody at 1:1000.

- 1- mpkCCD cell lysate (mouse)
- 2- FRT expressing tagged β -mENaC

Please check the product details page for more images. Overall 4 images are available for ABIN2486401.