
# antibodies -online.com





## anti-SLC9A3 antibody (PerCP)

2 Images



Go to Product page

#### Overview

Quantity:	100 μg
Target:	SLC9A3
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This SLC9A3 antibody is conjugated to PerCP
Application:	Western Blotting (WB), Immunocytochemistry (ICC), Immunohistochemistry (IHC), Immunofluorescence (IF)

#### **Product Details**

Immunogen:	Synthetic peptide mapping to AA 809 to 831 of rat sequence
Specificity:	Most abundant in colon and small intestine, followed by kidney and stomach. In kidney, expressed in proximal tubules and outer medulla (at protein level),Detects ~84 kDa.
Cross-Reactivity:	Mouse, Rat
Purification:	Protein A Purified

## Target Details

Target:	SLC9A3
Alternative Name:	NHE3 (SLC9A3 Products)
Background:	Sodium-hydrogen exchanger 3 (NHE3, Slc9a3) is an epithelial transport protein that carries out
	1:1 exchange of Na+ and H+ across the plasma membrane. It is apically located in the proximal

## **Target Details**

	tubule of the kidney, the thick ascending limb of the kidney, and in small intestine (1). NHE3 is phosphorylated and regulated by multiple kinases including PKA, SGK1 and CK2. It can be phosphorylated by calyculin A, and dephosphorylated by PP1 catalytic subunit in vitro (2).
Gene ID:	24784
NCBI Accession:	NP_036786
UniProt:	P26433
Pathways:	Proton Transport

## **Application Details**


Application Notes:	<ul> <li>WB (1:1000)</li> <li>IHC (1:25)</li> <li>optimal dilutions for assays should be determined by the user.</li> </ul>
Comment:	1 $\mu$ g/ml of ABIN2486550 was sufficient for detection of HNE3 in 10 $\mu$ g of rat kidney tissue lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

For Research Use only


## Handling

Restrictions:

Format:	Liquid
Concentration:	1 mg/mL
Buffer:	PBS, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Conjugated antibodies should be stored at 4°C



### L546 (NHE3) Immunoblot



#### **Immunohistochemistry**

Image 1. Immunohistochemistry analysis using Rabbit Anti-NHE3 Polyclonal Antibody (ABIN2486550). Tissue: Colon. Species: Rat. Fixation: Formalin Fixed Paraffin-Embedded. Primary Antibody: Rabbit Anti-NHE3 Polyclonal Antibody (ABIN2486550) at 1:25 for 1 hour at RT. Secondary Antibody: Goat Anti-Rabbit IgG:Alexa Fluor 488. Counterstain: Actin-binding Phalloidin-Alexa Fluor 633, DAPI (blue) nuclear stain. Magnification: 63X. (A) DAPI (blue) nuclear stain. (B) Phalloidin Alex Fluor 633 F-Actin stain. (C) NHE3 Antibody (D) Composite

#### **Western Blotting**

**Image 2.** Western blot analysis of Rat kidney tissue lysates showing detection of NHE3 protein using Rabbit Anti-NHE3 Polyclonal Antibody . Primary Antibody: Rabbit Anti-NHE3 Polyclonal Antibody at 1:1000.