



[Go to Product page](#)

Datasheet for ABIN2486706
anti-HSP90AB1 antibody (APC)

5 Images

Overview

| | |
|--------------|--|
| Quantity: | 100 µL |
| Target: | HSP90AB1 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This HSP90AB1 antibody is conjugated to APC |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), ELISA, Immunofluorescence (IF), Immunocytochemistry (ICC) |

Product Details

| | |
|-------------------|---|
| Immunogen: | Full length protein HSP90 |
| Specificity: | Detects ~90kda. Does not cross-react with HSP90α. |
| Cross-Reactivity: | Human, Mouse, Rat |
| Purification: | Peptide Affinity Purified |

Target Details

| | |
|-------------------|---|
| Target: | HSP90AB1 |
| Alternative Name: | HSP90 beta (HSP90AB1 Products) |
| Background: | HSP90 is a highly conserved and essential stress protein that is expressed in all eukaryotic cells. From a functional perspective, HSP90 participates in the folding, assembly, maturation, and stabilization of specific proteins as an integral component of a chaperone complex (1-4). |

Target Details

Despite its label of being a heat-shock protein, HSP90 is one of the most highly expressed proteins in unstressed cells (1-2 % of cytosolic protein). It carries out a number of housekeeping functions - including controlling the activity, turnover, and trafficking of a variety of proteins. Most of the HSP90-regulated proteins that have been discovered to date are involved in cell signaling (5-6). The number of proteins now known to interact with HSP90 is about 100. Target proteins include the kinases v-Src, Wee1, and c-Raf, transcriptional regulators such as p53 and steroid receptors, and the polymerases of the hepatitis B virus and telomerase.5. When bound to ATP, HSP90 interacts with co-chaperones Cdc37, p23, and an assortment of immunophilin-like proteins, forming a complex that stabilizes and protects target proteins from proteasomal degradation. In most cases, HSP90-interacting proteins have been shown to co-precipitate with HSP90 when carrying out immunoadsorption studies, and to exist in cytosolic heterocomplexes with it. In a number of cases, variations in HSP90 expression or HSP90 mutation has been shown to degrade signaling function via the protein or to impair a specific function of the protein (such as steroid binding, kinase activity) in vivo. Ansamycin antibiotics, such as geldanamycin and radicicol, inhibit HSP90 function (7). Looking for more information on HSP90? Visit our new HSP90 Scientific Resource Guide at <http://www.HSP90.ca>.

| | |
|-----------------|---|
| Gene ID: | 3326 |
| NCBI Accession: | NP_031381 |
| UniProt: | P08238 |
| Pathways: | Regulation of Cell Size |

Application Details

| | |
|--------------------|---|
| Application Notes: | <ul style="list-style-type: none">• WB (1:1000)• IHC (1:100)• ICC/IF (1:120)• optimal dilutions for assays should be determined by the user. |
| Comment: | A 1:1000 dilution of ABIN2486706 was sufficient for detection of HSP90 in 20 µg of HeLa cell lysate by ECL immunoblot analysis. |
| Restrictions: | For Research Use only |


Handling

| | |
|----------------|---------|
| Format: | Liquid |
| Concentration: | 1 mg/mL |

Handling


| | |
|--------------------|--|
| Buffer: | PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated |
| Preservative: | Sodium azide |
| Precaution of Use: | This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only. |
| Storage: | 4 °C |
| Storage Comment: | Conjugated antibodies should be stored at 4°C |

Images




Immunofluorescence (fixed cells)

Image 1. Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 beta Polyclonal Antibody . Tissue: HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 beta Polyclonal Antibody at 1:120 for 12 hours at 4°C. Secondary Antibody: R-PE Goat Anti-Rabbit (yellow) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Melanosome. Magnification: 20x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 beta Antibody. (C) Composite.



Western Blotting

Image 2. Western blot analysis of Human Cell line lysates showing detection of HSP90 beta protein using Rabbit Anti-HSP90 beta Polyclonal Antibody . Load: 15 µg protein. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-HSP90 beta Polyclonal Antibody at 1:2000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.



Immunofluorescence (fixed cells)

Image 3. Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-Hsp90 beta Polyclonal Antibody . Tissue: HeLa Cells. Species: Human. Fixation: 2% Formaldehyde for 20 min at RT. Primary Antibody: Rabbit Anti-Hsp90 beta Polyclonal Antibody at 1:120 for 12 hours at 4°C. Secondary Antibody: APC Goat Anti-Rabbit (red) at 1:200 for 2 hours at RT. Counterstain: DAPI (blue) nuclear stain at 1:40000 for 2 hours at RT. Localization: Cytoplasm. Melanosome. Magnification: 100x. (A) DAPI (blue) nuclear stain. (B) Anti-Hsp90 beta Antibody. (C) Composite.

Please check the [product details page](#) for more images. Overall 5 images are available for ABIN2486706.