

Datasheet for ABIN2486746  
**anti-HSPB8 antibody (PE)**[Go to Product page](#)

## 4 Images

## Overview

|              |  |
|--------------|--|
| Quantity:    | 100 µL   |
| Target:      | HSPB8  |
| Reactivity:  | Human  |
| Host:        | Rabbit   |
| Clonality:   | Polyclonal   |
| Conjugate:   | This HSPB8 antibody is conjugated to PE  |
| Application: | Western Blotting (WB), Immunohistochemistry (IHC), Immunofluorescence (IF),<br>Immunocytochemistry (ICC), Immunoprecipitation (IP) |

## Product Details

|                   |   |
|-------------------|---|
| Immunogen:        | Human HSP22   |
| Specificity:      | Detects ~22 kDa. Does not cross-react with HSP27 or alpha-crystallin. |
| Cross-Reactivity: | Human, Mouse, Rat   |
| Purification:     | Peptide Affinity Purified   |

## Target Details

|                   |  |
|-------------------|--|
| Target:           | HSPB8  |
| Alternative Name: | HSP22 ( <a href="#">HSPB8 Products</a> )   |
| Background:       | HSP22 (HSPB8) is a 196-amino acid protein that is a member of the small heat shock protein super-family and the human protein is most closely related to HSP27. Similar to most other sHSPs, HSP22 is predominately transcribed in skeletal muscle and heart, as well as the |

## Target Details

placenta (1). HSP22 is a monomeric protein which interacts with HSPB1. It displays temperature-dependent chaperone activity. In a two hybrid screen, HSPB8 interacted preferentially with a triple aspartate form of HSP27 which mimics HSP27 phosphorylated at Ser15, Ser78, and Ser82, as compared to wild-type HSP27 (2). HSPB8 has two binding domains (N and C Terminal) that are specific for different binding partners, and has the ability to bind itself and other sHSPs (3). The chaperone-like activity is of great importance to the function of HSP22 in various processes including proliferation, apoptosis and macro autophagy (4). Mutations in the HSPB8 gene are associated with the inherited peripheral neuropathies, autosomal dominant distal hereditary motor neuropathy type IIA (dSMA) and axonal Charcot-Marie-Tooth disease type 2L (CMT2L) (5).

Gene ID: 26353

NCBI Accession: [NP\\_055180](#)

UniProt: [Q9UJY1](#)

## Application Details

Application Notes:

- WB (1:1000)
- ICC/IF (1:100)
- IHC (1:100)
- optimal dilutions for assays should be determined by the user.

Comment: A 1:1000 dilution of ABIN2486746 was sufficient for detection of HSP22 in 10 µg of rat tissue lysate by colorimetric immunoblot analysis using Goat anti-rabbit IgG:HRP as the secondary antibody.

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 mg/mL

Buffer: PBS pH 7.4, 50 % glycerol, 0.09 % sodium azide, Storage buffer may change when conjugated


Preservative: Sodium azide

Precaution of Use: This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.

Storage: 4 °C


Storage Comment: Conjugated antibodies should be stored at 4°C

## Images




## Immunohistochemistry

**Image 1.** Immunohistochemistry analysis using Rabbit Anti-HSP22 Polyclonal Antibody . Tissue: Spinal cord. Species: Mouse. Primary Antibody: Rabbit Anti-HSP22 Polyclonal Antibody at 1:100. Secondary Antibody: Alexa Fluor 488 Goat Anti-Rabbit. DAPI merged with Alexa 488. Courtesy of: Joy Irobi-Devolder, University of Antwerp, Belgium..



## Immunofluorescence (fixed cells)

**Image 2.** Immunocytochemistry/Immunofluorescence analysis using Rabbit Anti-HSP22 Polyclonal Antibody . Tissue: HaCaT cells. Species: Human. Fixation: Cold 100% methanol at -20C for 10 minutes. Primary Antibody: Rabbit Anti-HSP22 Polyclonal Antibody at 1:100 for 12 hours at 4°C. Secondary Antibody: FITC Goat Anti-Rabbit at 1:50 for 1-2 hours at RT in dark. Localization: Nuclear Staining.



## Western Blotting

**Image 3.** Western blot analysis of Rat Skeletal muscle lysates showing detection of HSP22 protein using Rabbit Anti-HSP22 Polyclonal Antibody . Load: 15 μg protein. Block: 1.5% BSA for 30 minutes at RT. Primary Antibody: Rabbit Anti-HSP22 Polyclonal Antibody at 1:1000 for 2 hours at RT. Secondary Antibody: Donkey Anti-Rabbit IgG: HRP for 1 hour at RT.

Please check the [product details page](#) for more images. Overall 4 images are available for ABIN2486746.