

Datasheet for ABIN2535421

IL17C ELISA Kit



Overview

Quantity:	96 tests
Target:	IL17C
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	7.81 pg/mL - 500 pg/mL
Minimum Detection Limit:	7.81 pg/mL
Application:	ELISA
Product Details	
Purpose:	Human Interleukin 17C ELISA Kit is a sandwich ELISA kit for use with Serum, plasma and other
Purpose:	Human Interleukin 17C ELISA Kit is a sandwich ELISA kit for use with Serum, plasma and other biological fluids. This assay has high sensitivity and excellent specificity for detection of
Purpose:	
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Interleukin 17C (IL17C)
Purpose: Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Interleukin 17C (IL17C) No significant cross-reactivity or interference between Interleukin 17C (IL17C) and analogues
	biological fluids. This assay has high sensitivity and excellent specificity for detection of Interleukin 17C (IL17C) No significant cross-reactivity or interference between Interleukin 17C (IL17C) and analogues was observed.
Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Interleukin 17C (IL17C) No significant cross-reactivity or interference between Interleukin 17C (IL17C) and analogues was observed. Plasma, Serum
Sample Type: Analytical Method:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Interleukin 17C (IL17C) No significant cross-reactivity or interference between Interleukin 17C (IL17C) and analogues was observed. Plasma, Serum Quantitative

Target Details

Expiry Date:

6 months

larget Details	
Target:	IL17C
Abstract:	IL17C Products
Pathways:	Cellular Response to Molecule of Bacterial Origin
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.