

Datasheet for ABIN2536346

NAD-ME ELISA Kit



Overview

Sensitivity:

Quantity:	96 tests
Target:	NAD-ME
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human ME2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other
Purpose:	Human ME2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic
Purpose:	•
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2)
Purpose: Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) No significant cross-reactivity or interference between Malic Enzyme 2, NADP+ Dependent,
	biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) No significant cross-reactivity or interference between Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) and analogues was observed.
Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) No significant cross-reactivity or interference between Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) and analogues was observed. Tissue Homogenate

< 0.054 ng/mL

Target Details

Target:	NAD-ME
Alternative Name:	Malic Enzyme 2, NADP+ Dependent, Mitochondrial (ME2) (NAD-ME Products)
Pathways:	Production of Molecular Mediator of Immune Response
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
Expiry Date:	6 months