

Datasheet for ABIN2537076

NGRN ELISA Kit



Overview

- Overview	
Quantity:	96 tests
Target:	NGRN
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.781 ng/mL - 50 ng/mL
Minimum Detection Limit:	0.781 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human Neugrin ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other
Purpose:	Human Neugrin ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other biological fluids. This assay has high sensitivity and excellent specificity for detection of
Purpose:	
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neugrin (NGRN)
Purpose:	biological fluids. This assay has high sensitivity and excellent specificity for detection of
Purpose: Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neugrin (NGRN) No significant cross-reactivity or interference between Neugrin (NGRN) and analogues was
	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neugrin (NGRN) No significant cross-reactivity or interference between Neugrin (NGRN) and analogues was observed.
Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neugrin (NGRN) No significant cross-reactivity or interference between Neugrin (NGRN) and analogues was observed. Tissue Homogenate
Sample Type: Analytical Method:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neugrin (NGRN) No significant cross-reactivity or interference between Neugrin (NGRN) and analogues was observed. Tissue Homogenate Quantitative

Target Details

GRN Products) ty of the kit is determined by the rate of activity loss. The loss rate is less
ty of the kit is determined by the rate of activity loss. The loss rate is less
ty of the kit is determined by the rate of activity loss. The loss rate is less
expiration date under appropriate storage conditions. To minimize
ations, operation procedures and lab conditions should be strictly controlled.
uggested that the whole assay is performed by the same user throughout.
tions: Optimal dilutions/concentrations should be determined by the end
philized
kit is determined by the rate of activity loss. The loss rate is less than 5%
n date under appropriate storage conditions minimize performance
ion procedures and lab conditions should be strictly controlled. It is also
that the whole assay is performed by the same user throughout.
nly
the kit according to the storage instruction in the kit's manual.