

Datasheet for ABIN2537116

NKA ELISA Kit



Overview

Quantity:	96 tests
Target:	NKA
Reactivity:	Human
Method Type:	Competition ELISA
Detection Range:	12.35 pg/mL - 1000 pg/mL
Minimum Detection Limit:	12.35 pg/mL
Application:	ELISA
Product Details	
Purpose:	Human Neurokinin A ELISA Kit is a competitive ELISA kit for use with Serum, plasma and other
. d. pode.	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was observed.
Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was
·	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was observed.
Sample Type:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was observed. Plasma, Serum
Sample Type: Analytical Method:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was observed. Plasma, Serum Quantitative
Sample Type: Analytical Method: Detection Method:	biological fluids. This assay has high sensitivity and excellent specificity for detection of Neurokinin A (NKA) No significant cross-reactivity or interference between Neurokinin A (NKA) and analogues was observed. Plasma, Serum Quantitative Colorimetric

Target Details

Target:	NKA
Alternative Name:	Neurokinin A (NKA) (NKA Products)
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less
	than 5 % within the expiration date under appropriate storage conditions. To minimize
	performance fluctuations, operation procedures and lab conditions should be strictly controlled.
	It is also strongly suggested that the whole assay is performed by the same user throughout.
	Recommended dilutions: Optimal dilutions/concentrations should be determined by the end
	user.
	Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5%
	within the expiration date under appropriate storage conditions minimize performance
	fluctuations, operation procedures and lab conditions should be strictly controlled. It is also
	strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Llandling	
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
Expiry Date:	6 months