

Datasheet for ABIN2538533

RIPK1 ELISA Kit



oo to . Toddot pago

()	ve	r\/i	۱۸/
\cup	V C	1 / 1	 v v

Quantity:	96 tests
Target:	RIPK1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human RIPK1 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other
	biological fluids. This assay has high sensitivity and excellent specificity for detection of
	Receptor Interacting Serine Threonine Kinase 1 (RIPK1)
	No significant cross-reactivity or interference between Receptor Interacting Serine Threonine
	Kinase 1 (RIPK1) and analogues was observed.
Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Receptor Interacting
	Serine Threonine Kinase 1 (RIPK1)
Sensitivity:	< 0.059 ng/mL

Target Details

RIPK1
Receptor Interacting Serine Threonine Kinase 1 (RIPK1) (RIPK1 Products)
NF-kappaB Signaling, Apoptosis, Caspase Cascade in Apoptosis, TLR Signaling, Activation of
Innate immune Response, Inositol Metabolic Process, Positive Regulation of Endopeptidase
Activity, Hepatitis C, Protein targeting to Nucleus, Toll-Like Receptors Cascades, Negative
Regulation of intrinsic apoptotic Signaling, SARS-CoV-2 Protein Interactome, Ubiquitin
Proteasome Pathway
Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less
than 5 % within the expiration date under appropriate storage conditions. To minimize
performance fluctuations, operation procedures and lab conditions should be strictly controlled
It is also strongly suggested that the whole assay is performed by the same user throughout.
Recommended dilutions: Optimal dilutions/concentrations should be determined by the end
user.
Standard Form: Lyophilized
The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5%
within the expiration date under appropriate storage conditions. To minimize performance
fluctuations, operation procedures and lab conditions should be strictly controlled. It is also
strongly suggested that the whole assay is performed by the same user throughout.
Pre-coated
For Research Use only
4 °C/-20 °C
Upon receipt, store the kit according to the storage instruction in the kit's manual.
6 months