

## Datasheet for ABIN2539190

## Stanniocalcin 2 ELISA Kit



## Overview

Quantity:	96 tests
Target:	Stanniocalcin 2 (STC2)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.25 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.25 pg/mL
Application:	ELISA
Product Details	
Purpose:	Human Stanniocalcin 2 ELISA Kit is a sandwich ELISA kit for use with Serum, plasma, tissue
	homogenates and other biological fluids. This assay has high sensitivity and excellent specificity for detection of Stanniocalcin 2 (STC2)  No significant cross-reactivity or interference between Stanniocalcin 2 (STC2) and analogues was observed.
Sample Type:	specificity for detection of Stanniocalcin 2 (STC2)  No significant cross-reactivity or interference between Stanniocalcin 2 (STC2) and analogues
Sample Type: Analytical Method:	specificity for detection of Stanniocalcin 2 (STC2)  No significant cross-reactivity or interference between Stanniocalcin 2 (STC2) and analogues was observed.
	specificity for detection of Stanniocalcin 2 (STC2)  No significant cross-reactivity or interference between Stanniocalcin 2 (STC2) and analogues was observed.  Plasma, Serum, Tissue Homogenate
Analytical Method:	specificity for detection of Stanniocalcin 2 (STC2)  No significant cross-reactivity or interference between Stanniocalcin 2 (STC2) and analogues was observed.  Plasma, Serum, Tissue Homogenate  Quantitative

## Target Details

- 3	
Target:	Stanniocalcin 2 (STC2)
Abstract:	STC2 Products
Pathways:	Hormone Activity, ER-Nucleus Signaling, SARS-CoV-2 Protein Interactome
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.  Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
Expiry Date:	6 months