

Datasheet for ABIN2539346

SNCG ELISA Kit



Go to Product page

_						
	V	\triangle	r۱	/1	\triangle	Λ/
	' V '		ΙV			v v

Quantity:	96 tests
Target:	SNCG
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	31.25 pg/mL - 2000 pg/mL
Minimum Detection Limit:	31.25 pg/mL
Application:	ELISA
Product Details	
Purpose:	Human Synuclein Gamma ELISA Kit is a sandwich ELISA kit for use with Serum, plasma, tissue
	homogenates, cell lysates and other biological fluids. This assay has high sensitivity and
	excellent specificity for detection of Synuclein Gamma (SNCg)
	No significant cross-reactivity or interference between Synuclein Gamma (SNCg) and
	analogues was observed.
Sample Type:	Cell Lysate, Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Synuclein Gamma
	(SNCg)
Sensitivity:	< 13.5 pg/mL

Target Details

Application Details Application Notes: Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly control It is also strongly suggested that the whole assay is performed by the same user throughon Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 50 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated	
Application Notes: Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 50 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly control It is also strongly suggested that the whole assay is performed by the same user throughout Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
performance fluctuations, operation procedures and lab conditions should be strictly control it is also strongly suggested that the whole assay is performed by the same user throughout Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 50 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	í
It is also strongly suggested that the whole assay is performed by the same user throughor Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
Recommended dilutions: Optimal dilutions/concentrations should be determined by the enuser. Standard Form: Lyophilized The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	lled.
User. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	t.
Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	I
Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 55 within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.	
strongly suggested that the whole assay is performed by the same user throughout.	
Plate: Pre-coated	
Restrictions: For Research Use only	
Handling	
Storage: 4 °C/-20 °C	
Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	
Expiry Date: 6 months	