

## Datasheet for ABIN2540710

## **ADAM17 ELISA Kit**



## Overview

Sensitivity:

Quantity:	96 tests
Target:	ADAM17
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human ADAM17 ELISA Kit is a sandwich ELISA kit for use with Serum, plasma, tissue
	homogenates, cell lysates, cell culture supernates and other biological fluids. This assay has
	high sensitivity and excellent specificity for detection of A Disintegrin And Metalloproteinase 17
	(ADAM17)
	No significant cross-reactivity or interference between A Disintegrin And Metalloproteinase 17
	(ADAM17) and analogues was observed.
Sample Type:	Cell Culture Supernatant, Cell Lysate, Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of A Disintegrin And
	Metalloproteinase 17 (ADAM17)
On maritim site or	. 0.054 p.s./pal

< 0.054 ng/mL

## **Target Details**

Expiry Date:

6 months

rarget Details	
Target:	ADAM17
Alternative Name:	A Disintegrin And Metalloproteinase 17 (ADAM17) (ADAM17 Products)
Gene ID:	6
UniProt:	P78536
Pathways:	Notch Signaling, EGFR Signaling Pathway, Neurotrophin Signaling Pathway, Response to Growth Hormone Stimulus
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user.  Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.