

Datasheet for ABIN2540800 ACTG2 ELISA Kit



Overview

| 0000000 | |
|--------------------------|---|
| Quantity: | 96 tests |
| Target: | ACTG2 |
| Reactivity: | Human |
| Method Type: | Sandwich ELISA |
| Detection Range: | 0.156 ng/mL - 10 ng/mL |
| Minimum Detection Limit: | 0.156 ng/mL |
| Application: | ELISA |
| Product Details | |
| Purpose: | Human ACTg2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other |
| | biological fluids. This assay has high sensitivity and excellent specificity for detection of Actin |
| | Gamma 2, Smooth Muscle (ACTg2) |
| | No significant cross-reactivity or interference between Actin Gamma 2, Smooth Muscle |
| | (ACTg2) and analogues was observed. |
| Sample Type: | Tissue Homogenate |
| Analytical Method: | Quantitative |
| Detection Method: | Colorimetric |
| Specificity: | This assay has high sensitivity and excellent specificity for detection of Actin Gamma 2, |
| | Smooth Muscle (ACTg2) |
| Sensitivity: | < 0.053 ng/mL |

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2540800 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

| Target Details | |
|---------------------|--|
| Target: | ACTG2 |
| Alternative Name: | Actin gamma 2, Smooth Muscle (ACTg2) (ACTG2 Products) |
| Application Details | |
| Application Notes: | Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized |
| Comment: | The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. |
| Plate: | Pre-coated |
| Restrictions: | For Research Use only |
| Handling | |
| Storage: | 4 °C/-20 °C |
| Storage Comment: | Upon receipt, store the kit according to the storage instruction in the kit's manual. |
| Expiry Date: | 6 months |