ANTIBODIES ONLINE

Datasheet for ABIN2541576 TAP1 ELISA Kit



Overview

Quantity:	96 tests
Target:	TAP1
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human TAP1 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates, cell lysates
	and other biological fluids. This assay has high sensitivity and excellent specificity for detection
	of Antigen Peptide Transporter 1 (TAP1)
	No significant cross-reactivity or interference between Antigen Peptide Transporter 1 (TAP1)
	and analogues was observed.
Sample Type:	Cell Lysate, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Antigen Peptide
	Transporter 1 (TAP1)
Sensitivity:	< 0.134 ng/mL

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2541576 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details

Target:	TAP1
Alternative Name:	Antigen Peptide Transporter 1 (TAP1) (TAP1 Products)
Pathways:	Regulation of Leukocyte Mediated Immunity, Positive Regulation of Immune Effector Process,
	Human Leukocyte Antigen (HLA) in Adaptive Immune Response

Application Details

Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize
performance fluctuations, operation procedures and lab conditions should be strictly controlled
It is also strongly suggested that the whole assay is performed by the same user throughout.
Recommended dilutions: Optimal dilutions/concentrations should be determined by the end
user.
Standard Form: Lyophilized
The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5%
within the expiration date under appropriate storage conditions minimize performance
fluctuations, operation procedures and lab conditions should be strictly controlled. It is also
strongly suggested that the whole assay is performed by the same user throughout.
Pre-coated
For Research Use only
4 °C/-20 °C
Upon receipt, store the kit according to the storage instruction in the kit's manual.
6 months