

Datasheet for ABIN2542827

CCT2 ELISA Kit



Overview	
Quantity:	96 tests
Target:	CCT2
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human CCT2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other
	biological fluids. This assay has high sensitivity and excellent specificity for detection of
	Chaperonin Containing TCP1, Subunit 2 (CCT2)
	No significant cross-reactivity or interference between Chaperonin Containing TCP1, Subunit 2

Product Details	
Purpose:	Human CCT2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and other biological fluids. This assay has high sensitivity and excellent specificity for detection of Chaperonin Containing TCP1, Subunit 2 (CCT2) No significant cross-reactivity or interference between Chaperonin Containing TCP1, Subunit 2 (CCT2) and analogues was observed.
Sample Type:	Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Chaperonin Containing TCP1, Subunit 2 (CCT2)
Sensitivity:	< 0.056 ng/mL

Target Details

Alternative Name: Chaperonin Containing TCP1, Subunit 2 (CCT2) (CCT2 Products) Application Details Application Notes: Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled it is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual. Expiry Date: 6 months	Target:	CCT2
Application Notes: Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Alternative Name:	Chaperonin Containing TCP1, Subunit 2 (CCT2) (CCT2 Products)
than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Application Details	
performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less
It is also strongly suggested that the whole assay is performed by the same user throughout. Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		than 5 % within the expiration date under appropriate storage conditions. To minimize
Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		performance fluctuations, operation procedures and lab conditions should be strictly controlled.
User. Standard Form: Lyophilized Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		It is also strongly suggested that the whole assay is performed by the same user throughout.
Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		Recommended dilutions: Optimal dilutions/concentrations should be determined by the end
Comment: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		user.
within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		Standard Form: Lyophilized
fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5%
strongly suggested that the whole assay is performed by the same user throughout. Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		within the expiration date under appropriate storage conditions. To minimize performance
Plate: Pre-coated Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		fluctuations, operation procedures and lab conditions should be strictly controlled. It is also
Restrictions: For Research Use only Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		strongly suggested that the whole assay is performed by the same user throughout.
Handling Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Plate:	Pre-coated
Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Restrictions:	For Research Use only
Storage: 4 °C/-20 °C Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.		
Storage Comment: Upon receipt, store the kit according to the storage instruction in the kit's manual.	Handling	
	Storage:	4 °C/-20 °C
Expiry Date: 6 months	Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
	Expiry Date:	6 months