

Datasheet for ABIN2543808

Desmocollin 2 ELISA Kit



Overview

Quantity:	96 tests
Target:	Desmocollin 2 (DSC2)
Reactivity:	Human
Method Type:	Sandwich ELISA
Detection Range:	0.312 ng/mL - 20 ng/mL
Minimum Detection Limit:	0.312 ng/mL
Application:	ELISA
Product Details	
Purpose:	Human Desmocollin 2 ELISA Kit is a sandwich ELISA kit for use with Tissue homogenates and
	other biological fluids. This assay has high sensitivity and excellent specificity for detection of Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues was observed.
Sample Type:	Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues
Sample Type: Analytical Method:	Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues was observed.
	Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues was observed. Tissue Homogenate
Analytical Method:	Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues was observed. Tissue Homogenate Quantitative
Analytical Method: Detection Method:	Desmocollin 2 (DSC2) No significant cross-reactivity or interference between Desmocollin 2 (DSC2) and analogues was observed. Tissue Homogenate Quantitative Colorimetric

Target Details

Target:	Desmocollin 2 (DSC2)
Abstract:	DSC2 Products
Application Details	
Application Notes:	Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5 % within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
	Recommended dilutions: Optimal dilutions/concentrations should be determined by the end user. Standard Form: Lyophilized
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
Expiry Date:	6 months