

## Datasheet for ABIN2543824 **DGAT1 ELISA Kit**



Overview

Quantity:	96 tests
Target:	DGAT1
Reactivity:	Mouse
Method Type:	Sandwich ELISA
Detection Range:	0.156 ng/mL - 10 ng/mL
Minimum Detection Limit:	0.156 ng/mL
Application:	ELISA
Product Details	
Purpose:	Mouse DGAT1 ELISA Kit is a sandwich ELISA kit for use with Serum, plasma, tissue
	homogenates and other biological fluids. This assay has high sensitivity and excellent
	specificity for detection of Diacylglycerol-O-Acyltransferase Homolog 1 (DGAT1)
	No significant cross-reactivity or interference between Diacylglycerol-O-Acyltransferase
	Homolog 1 (DGAT1) and analogues was observed.
Sample Type:	Plasma, Serum, Tissue Homogenate
Analytical Method:	Quantitative
Detection Method:	Colorimetric
Specificity:	This assay has high sensitivity and excellent specificity for detection of Diacylglycerol-O-Acy
	Itransferase Homolog 1 (DGAT1)
Sensitivity:	< 0.054 ng/mL

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/2 | Product datasheet for ABIN2543824 | 07/26/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Target:	DGAT1
Alternative Name:	Diacylglycerol-O-Acyltransferase Homolog 1 (DGAT1) (DGAT1 Products)
Pathways:	Hormone Transport
Application Details	
Application Notes:	<ul> <li>Stability: The stability of the kit is determined by the rate of activity loss. The loss rate is less</li> <li>than 5 % within the expiration date under appropriate storage conditions. To minimize</li> <li>performance fluctuations, operation procedures and lab conditions should be strictly controlled.</li> <li>It is also strongly suggested that the whole assay is performed by the same user throughout.</li> <li>Recommended dilutions: Optimal dilutions/concentrations should be determined by the end</li> <li>user.</li> <li>Standard Form: Lyophilized</li> </ul>
Comment:	The stability of the kit is determined by the rate of activity loss. The loss rate is less than 5% within the expiration date under appropriate storage conditions. To minimize performance fluctuations, operation procedures and lab conditions should be strictly controlled. It is also strongly suggested that the whole assay is performed by the same user throughout.
Plate:	Pre-coated
Restrictions:	For Research Use only
Handling	
Storage:	4 °C/-20 °C
Storage Comment:	Upon receipt, store the kit according to the storage instruction in the kit's manual.
Expiry Date:	6 months