antibodies -online.com





anti-GRIP1 antibody (AA 1-100) (Biotin)



Overview

| Quantity: | 100 μL |
|----------------------|---|
| Target: | GRIP1 |
| Binding Specificity: | AA 1-100 |
| Reactivity: | Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This GRIP1 antibody is conjugated to Biotin |
| Application: | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human GRIP1 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Mouse |
| Predicted Reactivity: | Human,Rat,Cow |
| Purification: | Purified by Protein A. |

Target Details

| Target: | GRIP1 |
|-------------------|------------------------|
| Alternative Name: | GRIP1 (GRIP1 Products) |

Target Details

Expiry Date:

12 months

| rarget Details | |
|---------------------|---|
| Background: | Synonyms: GRIP, Glutamate receptor-interacting protein 1, GRIP-1, GRIP1 Background: May play a role as a localized scaffold for the assembly of a multiprotein signaling complex and as mediator of the trafficking of its binding partners at specific subcellular location in neurons. |
| Gene ID: | 23426 |
| UniProt: | Q9Y3R0 |
| Pathways: | Intracellular Steroid Hormone Receptor Signaling Pathway |
| Application Details | |
| Application Notes: | WB 1:300-5000 |
| | IHC-P 1:200-400 |
| | IHC-F 1:100-500 |
| Restrictions: | For Research Use only |
| Handling | |
| Format: | Liquid |
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C for 12 months. |
| Evein Deter | 10 months |