

## Datasheet for ABIN2559516 anti-IVD antibody (AA 201-300) (HRP)



Overview	
Quantity:	100 μL
Target:	IVD
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IVD antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human IVD
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Dog,Cow,Sheep,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	IVD

## **Target Details**

Background:	Synonyms: ACAD2, Isovaleryl-CoA dehydrogenase, mitochondrial, IVD
	Background: Isovaleryl-CoA dehydrogenase (IVD) is a mitochondrial matrix enzyme that
	catalyzes the third step in leucine catabolism. The genetic deficiency of IVD results in an
	accumulation of isovaleric acid, which is toxic to the central nervous system and leads to
	isovaleric acidemia. Alternatively spliced transcript variants encoding different isoforms have
	been found for this gene. [provided by RefSeq, Apr 2009]
Gene ID:	3712
UniProt:	P26440
Pathways:	Monocarboxylic Acid Catabolic Process
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months