

[Go to Product page](#)

Datasheet for ABIN2559621  
**anti-EIF3G antibody (AA 51-150) (Biotin)**

## Overview

Quantity:	100 µL
Target:	EIF3G
Binding Specificity:	AA 51-150
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EIF3G antibody is conjugated to Biotin
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human EIF3S4
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Pig,Horse
Purification:	Purified by Protein A.

## Target Details

Target:	EIF3G
Alternative Name:	EIF3S4 ( <a href="#">EIF3G Products</a> )

## Target Details

Background:	<p>Synonyms: EIF3S4, EIF3-P42, eIF3-p44, eIF3-delta, Eukaryotic translation initiation factor 3 subunit G, eIF3g, Eukaryotic translation initiation factor 3 RNA-binding subunit, eIF-3 RNA-binding subunit, Eukaryotic translation initiation factor 3 subunit 4, eIF-3-delta, eIF3 p42, eIF3 p44</p> <p>Background: Component of the eukaryotic translation initiation factor 3 (eIF-3) complex, which is required for several steps in the initiation of protein synthesis. The eIF-3 complex associates with the 40S ribosome and facilitates the recruitment of eIF-1, eIF-1A, eIF-2:GTP:methionyl-tRNAi and eIF-5 to form the 43S preinitiation complex (43S PIC). The eIF-3 complex stimulates mRNA recruitment to the 43S PIC and scanning of the mRNA for AUG recognition. The eIF-3 complex is also required for disassembly and recycling of post-termination ribosomal complexes and subsequently prevents premature joining of the 40S and 60S ribosomal subunits prior to initiation. This subunit can bind 18S rRNA.</p>
Gene ID:	8666
UniProt:	<a href="#">O75821</a>
Pathways:	<a href="#">Ribonucleoprotein Complex Subunit Organization</a>

## Application Details

Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

---

Storage Comment:                      Store at -20°C for 12 months.

---

Expiry Date:                              12 months