

### Datasheet for ABIN2567158

# anti-CD11c antibody



_					
	W	0	rv	10	W

Quantity:	0.1 mg	
Target:	CD11c (ITGAX)	
Reactivity:	Mouse	
Host:	Hamster	
Clonality:	Monoclonal	
Conjugate:	This CD11c antibody is un-conjugated	
Application:	Flow Cytometry (FACS), Immunohistochemistry (IHC), Immunofluorescence (IF), Immunoprecipitation (IP), Functional Studies (Func)	

### **Product Details**

Clone:	N418	
Isotype:	IgG	
Specificity:	The N418 monoclonal antibody specifically reacts with the integrin $\alpha x$ chain of the mouse CD11c, which is expressed on dendritic cells, CD4-/CD8+ intestinal intraepithelial lymphocytes (IEL) and some activated T lymphocytes.	
Purification:	The monoclonal antibody was purified utilizing affinity chromatography and unreacted dye was removed from the product.	

#### **Target Details**

Target:	CD11c (ITGAX)
Alternative Name:	CD11c (ITGAX Products)

## **Target Details**

3		
Background:	The N418 monoclonal antibody specifically reacts with the integrin ax chain of the mouse CD11c, which is expressed on dendritic cells, CD4-/CD8+ intestinal intraepithelial lymphocytes (IEL) and some activated T lymphocytes. Low levels of CD11c were detected on mouse splenic natural killer cells and on the monocyte/macrophage lineage cells.CD11c expression is upregulated on IEL and T lymphocytes after activation. It binds to CD54, fibrinogen and iC3b and influences the leukocyte adhesive interactions. The N418 antibody binds to CD11c on splenic dendritic cells of the mouse in the T-dependent areas. It also contributes to the binding of iC3b.	
Gene ID:	16411	
UniProt:	Q9QXH4	
Pathways:	Complement System, Activated T Cell Proliferation, Integrin Complex	
Application Details		
Application Notes:	Optimal working dilution should be determined by the investigator.	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	0.5 mg/mL	
Buffer:	Phosphate-buffered aqueous solution, ≤0.09 % Sodium azide, may contain carrier protein/stabilizer, ph7.2.	
Storage:	4 °C	
Storage Comment:	The product should be stored undiluted at 4°C and should be protected from prolonged exposure to light. Do not freeze.	