

Datasheet for ABIN2613475
anti-SCARB1 antibody (AA 238-250)[Go to Product page](#)

2 Images

Overview

Quantity:	100 µg
Target:	SCARB1
Binding Specificity:	AA 238-250
Reactivity:	Human
Host:	Goat
Clonality:	Polyclonal
Conjugate:	This SCARB1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (IHC)

Product Details

Purpose:	SCARB1 / SR-BI (aa238-250)
Sequence:	NGLSKVDFWH SDQ
Isotype:	IgG
Specificity:	This antibody is expected to recognize both reported isoforms (NP_005496.4, NP_001076428.1).
Cross-Reactivity:	Human
Purification:	Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.
Grade:	Verified

Target Details


Target:	SCARB1
Alternative Name:	SCARB1 (SCARB1 Products)
Background:	SCARB1, scavenger receptor class B, member 1, CD36L1, CLA-1, CLA1, HDLQTL6, SR-BI, SRB1, CD36 and LIMPII analogous 1, CD36 antigen (collagen type I receptor, thrombospondin receptor)-like 1, scavenger receptor class B member 1, scavenger receptor class B
Gene ID:	949
NCBI Accession:	NP_005496 , NP_001076428
Pathways:	Cellular Response to Molecule of Bacterial Origin , Hepatitis C , Lipid Metabolism , SARS-CoV-2 Protein Interactome

Application Details

Application Notes:	Immunohistochemistry: Paraffin embedded Human Testis. Recommended concentration: 5 µg/mL. Western Blot: Approx 85 kDa band observed in Human Adrenal gland lysates (calculated MW of 57.0 kDa according to NP_005496.4). Recommended concentration: 0.01-0.03 µg/mL. Peptide ELISA: antibody detection limit dilution 1:128000.
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	0.5 mg/mL
Buffer:	Supplied at 0.5 mg/mL in Tris saline, 0.02 % sodium azide, pH 7.3 with 0.5 % bovine serum albumin.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Handling Advice:	Minimize freezing and thawing.
Storage:	-20 °C
Storage Comment:	Aliquot and store at -20°C, with minimal freeze/thawing. A working aliquot may be refrigerated at 4°C for a few weeks and still remain viable.



Immunohistochemistry

Image 1. Staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



Western Blotting

Image 2. ABIN2613475 (0.03µg/ml) staining of Human Adrenal Gland lysate (35µg protein in RIPA buffer). Primary incubation was 1 hour. Detected by chemiluminescence.