

Datasheet for ABIN2617917

anti-Caspase 1 antibody (AA 135-152)[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	Caspase 1 (CASP1)
Binding Specificity:	AA 135-152
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Caspase 1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Immunogen:	A synthetic peptide corresponding to a sequence at the N-terminus of human CASP1(135-152aa CSLEEAQRIWKQKSAEIY).
Isotype:	IgG
Specificity:	Expressed in larger amounts in spleen and lung. Detected in liver, heart, small intestine, colon, thymus, prostate, skeletal muscle, peripheral blood leukocytes, kidney and testis. No expression in the brain. .
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Purification:	Immunogen affinity purified

Target Details

Target:	Caspase 1 (CASP1)
Alternative Name:	CASP1 / Caspase 1 (CASP1 Products)
Background:	Name/Gene ID: CASP1 Subfamily: Cysteine C14 Family: Protease Synonyms: CASP1, Caspase-1, CASP-1, CASP1 nirs variant 1, Caspase 1, ICE, ICE-like protease, IL-1 beta-converting enzyme, IL-1BC, IL1BC, IL1BCE, p45, IL1B-convertase, Interleukin-1 beta convertase
Gene ID:	834
Pathways:	Apoptosis , Interferon-gamma Pathway , Positive Regulation of Endopeptidase Activity , Inflammasome

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , 0.05 mg Thimerosal, 0.05 mg sodium azide per 100 µg antibody.
Preservative:	Sodium azide, Thimerosal (Merthiolate)
Precaution of Use:	This product contains Sodium azide and Thimerosal (Merthiolate): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for 1 year. After reconstitution, at 4°C for 1 month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid freeze-thaw cycles.