

Datasheet for ABIN2620359
anti-MUC1 antibody (AA 1067-1083)



[Go to Product page](#)

Overview

Quantity:	100 µg
Target:	MUC1
Binding Specificity:	AA 1067-1083
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MUC1 antibody is un-conjugated
Application:	Western Blotting (WB)

Product Details

Immunogen:	A synthetic peptide corresponding to a sequence at the C-terminus of human MUC1(1067-1083aa QELQRDISEMFLQIYKQ).
Isotype:	IgG
Specificity:	Expressed on the apical surface of epithelial cells, especially of airway passages, breast and uterus. Also expressed in activated and unactivated T-cells. Overexpressed in epithelial tumors, such as breast or ovarian cancer and also in non-epithelial tumor cells. Isoform 7 is expressed in tumor cells only. .
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Purification:	Immunogen affinity purified

Target Details

Target:	MUC1
Alternative Name:	EMA / MUC1 (MUC1 Products)
Background:	Name/Gene ID: MUC1 Family: Mucin Synonyms: MUC1, CA15-3, CD227, Carcinoma-associated mucin, CD227 antigen, DF3 antigen, Episialin, Epithelial membrane antigen, H23 antigen, H23AG, Krebs von den Lungen-6, MAM6, MUC1/ZD, MUC-1/SEC, Mucin-1, Pem, KL-6, MUC-1/X, Tumor-associated mucin, Peanut-reactive urinary mucin, Polymorphic epithelial mucin, PUM, EMA, MUC-1, Mucin 1, transmembrane
Gene ID:	4582
Pathways:	Negative Regulation of intrinsic apoptotic Signaling

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na ₂ HPO ₄ , 0.05 mg Thimerosal, 0.05 mg sodium azide per 100 µg antibody.
Preservative:	Sodium azide, Thimerosal (Merthiolate)
Precaution of Use:	This product contains Sodium azide and Thimerosal (Merthiolate): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C, -20 °C
Storage Comment:	At -20°C for 1 year. After reconstitution, at 4°C for 1 month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid freeze-thaw cycles.