

Datasheet for ABIN2621751 anti-HDAC3 antibody (AA 1-18)



Overview	
Quantity:	100 μg
Target:	HDAC3
Binding Specificity:	AA 1-18
Reactivity:	Human, Mouse, Rat, Pig, Rabbit, Chicken, Guinea Pig, Horse, Monkey, Bat, Xenopus laevis, Sheep
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HDAC3 antibody is un-conjugated
Application:	Western Blotting (WB), Immunohistochemistry (IHC), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))
Product Details	
Immunogen:	A synthetic peptide corresponding to a sequence at the C-terminus of human HDAC3(411-428aa NEFYDGDHDNDKESDVEI), identical to the related rat and mouse sequences.
Isotype:	IgG
Specificity:	Widely expressed.
Cross-Reactivity (Details):	No cross reactivity with other proteins.
Purification:	Immunogen affinity purified
Target Details	
Target:	HDAC3

Target Details

Alternative Name:	HDAC3 (HDAC3 Products)
Background:	Name/Gene ID: HDAC3 Family: Histone Deacetylase
	Synonyms: HDAC3, RPD3, SMAP45, HD3, Histone deacetylase 3, RPD3-2
Gene ID:	8841
Pathways:	Neurotrophin Signaling Pathway, Regulation of Lipid Metabolism by PPARalpha, Regulation of Muscle Cell Differentiation, Skeletal Muscle Fiber Development

Application Details

Application Notes:	Optimal working dilution should be determined by the investigator.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Reconstitution:	Distilled water
Concentration:	Lot specific
Buffer:	Contains 5 mg BSA, 0.9 mg NaCl, 0.2 mg Na2 HPO4, 0.05 mg Thimerosal, 0.05 mg sodium azide per 100 µg antibody.
Preservative:	Sodium azide, Thimerosal (Merthiolate)
Precaution of Use:	This product contains Sodium azide and Thimerosal (Merthiolate): POISONOUS AND HAZARDOUS SUBSTANCES which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	At -20°C for 1 year. After reconstitution, at 4°C for 1 month. It can also be aliquotted and stored frozen at -20°C for a longer time. Avoid freeze-thaw cycles.