

Datasheet for ABIN2621979
anti-HNRNPH1 antibody (C-Term)[Go to Product page](#)

1 Image

Overview

Quantity:	50 µL
Target:	HNRNPH1
Binding Specificity:	AA 400-448, C-Term
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HNRNPH1 antibody is un-conjugated
Application:	Immunohistochemistry (IHC), Immunofluorescence (IF), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))

Product Details

Brand:	IHC-plus™
Isotype:	IgG
Specificity:	Region between residue 400 and the C-terminus (residue 448) of human Heterogeneous Nuclear Ribonucleoprotein H1 using the numbering given in entry NP_005511.1 (GeneID 3187).
Purification:	Immunoaffinity purified

Target Details

Target:	HNRNPH1
Alternative Name:	HNRNPH1 / HnRNP H (HNRNPH1 Products)

Target Details


Background:	Name/Gene ID: HNRNPH1
	Synonyms: HNRNPH1, HNRPH1, HnRNPH, HnRNP H, HNRPH
Gene ID:	3187

Application Details

Application Notes:	Approved: IF (1:50 - 1:500), IHC, IHC-P (1:100)
	Usage: Immunohistochemistry: Antigen retrieval is recommended. Antigen retrieval with citrate buffer will enhance staining. Likely to work with frozen sections. In some cases, the antibody may be diluted further than indicated. Human controls: Breast Carcinoma, Colon Carcinoma, Linitis Plastica Stomach Cancer, Non-Small Cell Lung Cancer, Ovarian Carcinoma, Pancreatic Islet Cell Tumor, Prostate Carcinoma, Skin Basal Cell Carcinoma, Skin Squamous Cell Carcinoma, Small Cell Lung Cancer, Stomach Adenocarcinoma. Mouse controls: Squamous Cell Carcinoma.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	Lot specific
Buffer:	Tris-buffered saline, 0.1 % BSA, 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C for up to 1 year.
Expiry Date:	12 months



Immunohistochemistry

Image 1. Anti-HNRNPH1 / HnRNP H antibody IHC staining of human tonsil. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 1:100.