# antibodies -online.com





# anti-TCERG1 antibody (AA 15-50)





Go to Product page

_					
	W	0	rv	10	W

Quantity:	50 μL	
Target:	TCERG1	
Binding Specificity:	AA 15-50	
Reactivity:	Human, Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TCERG1 antibody is un-conjugated	
Application:	Immunohistochemistry (IHC), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (IF)	

#### **Product Details**


Brand:	IHC-plus™
Isotype:	IgG
Specificity:	Region between residue 15 and 50 of human APPLICATIONS Transcription Factor CA150 (Transcription Elongation Regulator 1) using the numbering given in entry NP_006697.2 (GenelD 10915).
Purification:	Immunoaffinity purified

## **Target Details**

Target:	TCERG1	
Alternative Name:	TCERG1 / CA150 (TCERG1 Products)	

### **Target Details**

rarget Details	
Background:	Name/Gene ID: TCERG1
	Synonyms: TCERG1, CA150, Co-activator of 150 kDa, TAF2S, Transcription factor CA150, Urn1
Gene ID:	10915
Application Details	
Application Notes:	Approved: IF (1:50 - 1:500), IHC, IHC-P (1:100)
	Usage: Immunohistochemistry: Antigen retrieval is recommended. Antigen retrieval with citrate buffer will enhance staining. Likely to work with frozen sections. In some cases, the antibody may be diluted further than indicated. Human controls: Breast Carcinoma, Colon Carcinoma, Prostate Carcinoma, Small Cell Lung Cancer. Mouse controls: Squamous Cell Carcinoma.
Comment:	Target Species of Antibody: Human
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	Lot specific
Buffer:	Tris-buffered saline, 0.1 % BSA, 0.09 % sodium azide.
Preservative:	Sodium azide
Precaution of Use:	This product contains Sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which should be handled by trained staff only.
Storage:	4 °C
Storage Comment:	Store at 2-8°C for up to 1 year.
Expiry Date:	12 months



#### **Immunohistochemistry**

**Image 1.** Anti-TCERG1 / CA150 antibody IHC staining of human skin. Immunohistochemistry of formalin-fixed, paraffin-embedded tissue after heat-induced antigen retrieval. Antibody dilution 1:100.